

COMPACTIFICATION OF STRATA OF ABELIAN DIFFERENTIALS

MATT BAINBRIDGE, DAWEI CHEN, QUENTIN GENDRON, SAMUEL GRUSHEVSKY,
AND MARTIN MÖLLER

ABSTRACT. We describe the closure of the strata of abelian differentials with prescribed type of zeros and poles, in the projectivized Hodge bundle over the Deligne-Mumford moduli space of stable curves with marked points. We provide an explicit characterization of pointed stable differentials in the boundary of the closure, both a complex analytic proof and a flat geometric proof for smoothing the boundary differentials, and numerous examples. The main new ingredient in our description is a global residue condition arising from a full order on the dual graph of a stable curve.

CONTENTS

1. Introduction	1
2. Comparison of strata compactifications	10
3. Examples of the incidence variety compactification	15
4. Plumbing and gluing differentials	33
5. Flat geometric smoothing	47
References	57

1. INTRODUCTION

1.1. **Background.** The Hodge bundle $\Omega\mathcal{M}_g$ is a complex vector bundle of rank g over the moduli space \mathcal{M}_g of genus g Riemann surfaces. A point $(X, \omega) \in \Omega\mathcal{M}_g$ consists of a Riemann surface X of genus g and a (holomorphic) abelian differential ω on X . The complement of the zero section $\Omega\mathcal{M}_g^*$ is naturally stratified into strata $\Omega\mathcal{M}_g(\mu)$ where the multiplicity of all the zeros of ω is prescribed by a partition $\mu = (m_1, \dots, m_n)$ of $2g - 2$. By scaling the differentials, \mathbb{C}^* acts on $\Omega\mathcal{M}_g^*$ and preserves the stratification, hence one can consider the projectivized strata $\mathbb{P}\Omega\mathcal{M}_g(\mu)$ in the projectivized Hodge bundle $\mathbb{P}\Omega\mathcal{M}_g = \Omega\mathcal{M}_g^*/\mathbb{C}^*$. An abelian differential ω defines a flat metric with conical

Date: March 30, 2018.

Research of the first author is supported in part by the Simons Foundation grant #359821.

Research of the second author is supported in part by the National Science Foundation under the CAREER grant DMS-13-50396 and a Boston College Research Incentive Grant.

Research of the third author was supported in part by ERC-StG 257137.

Research of the fourth author is supported in part by the National Science Foundation under the grants DMS-12-01369 and DMS-15-01265, and by a Simons Fellowship in Mathematics (Simons Foundation grant #341858 to Samuel Grushevsky).

Research of the fifth author is supported in part by ERC-StG 257137.

singularities such that the underlying Riemann surface X can be realized as a plane polygon whose edges are pairwise identified via translation. In this sense (X, ω) is called a flat surface or a translation surface. Varying the shape of flat surfaces induces a $\mathrm{GL}_2^+(\mathbb{R})$ -action on the strata of abelian differentials, called Teichmüller dynamics. A number of questions about surface geometry boil down to understanding the $\mathrm{GL}_2^+(\mathbb{R})$ -orbit closures in Teichmüller dynamics. What are their dimensions? Do they possess manifold structures? How can one calculate relevant dynamical invariants? From the viewpoint of algebraic geometry the orbit closures are of an independent interest for cycle class computations, which can provide crucial information for understanding the geometry of moduli spaces.

Many of these questions can be better accessed if one can describe a geometrically meaningful compactification of the strata. In particular, the recent breakthrough of Eskin, Mirzakhani, Mohammadi [EM13; EMM15] and Filip [Fil16] shows that any orbit closure (under the standard topology) is a quasiprojective subvariety of a stratum. Thus describing the projective subvarieties that are closures of orbits in a compactified stratum can shed further light on the classification of orbit closures.

Identify Riemann surfaces with smooth complex curves. The Deligne-Mumford compactification $\overline{\mathcal{M}}_g$ of \mathcal{M}_g parameterizes stable genus g curves that are (at worst) nodal curves with finite automorphism groups. The Hodge bundle $\Omega\mathcal{M}_g$ extends as a rank g complex vector bundle $\Omega\overline{\mathcal{M}}_g$ over $\overline{\mathcal{M}}_g$. The fiber of $\Omega\overline{\mathcal{M}}_g$ over a nodal curve X parameterizes stable differentials that have (at worst) simple poles at the nodes of X with opposite residues on the two branches of a node. One way of compactifying $\mathbb{P}\Omega\mathcal{M}_g(\mu)$ is by taking its closure in the projectivized Hodge bundle $\mathbb{P}\Omega\overline{\mathcal{M}}_g$ over $\overline{\mathcal{M}}_g$, and we call it the *Hodge bundle compactification* of the strata.

Alternatively, one can lift a stratum $\mathbb{P}\Omega\mathcal{M}_g(\mu)$ to the moduli space $\mathcal{M}_{g,n}$ of genus g curves with n marked points by adding on each curve the data of the zeroes of differentials. Let $\overline{\mathcal{M}}_{g,n}$ be the Deligne-Mumford compactification of $\mathcal{M}_{g,n}$ that parameterizes stable genus g curves with n marked points. Taking the closure of $\mathbb{P}\Omega\mathcal{M}_g(\mu)$ in $\overline{\mathcal{M}}_{g,n}$ provides another compactification, which we call the *Deligne-Mumford compactification* of the strata.

By combining the two viewpoints above, in this paper we describe a strata compactification that we call the *incidence variety compactification* $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,n}^{\mathrm{inc}}(\mu)$. Let $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,n}$ be the projectivized Hodge bundle over $\overline{\mathcal{M}}_{g,n}$, which parameterizes *pointed stable differentials*. Then the incidence variety compactification of $\mathbb{P}\Omega\mathcal{M}_g(\mu)$ is defined as the closure of the stratum in the projectivization $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,n}$.

The incidence variety compactification records both the limit stable differentials and the limit positions of the zeros when abelian differentials become identically zero on some irreducible components of the nodal curve. It contains more information than the Hodge bundle compactification, because the latter loses the information about the limit positions of the zeros on the components of nodal curves where the stable differentials vanish identically. The incidence variety compactification also contains more information than the Deligne-Mumford compactification, because the latter loses the information on the relative sizes of flat surfaces corresponding to the components of nodal curves where the stable differentials are not identically zero.

Our characterization of the boundary of the incidence variety compactification is in terms of a collection of (possibly meromorphic) differentials on the components of a pointed stable curve that satisfy certain combinatorial and residue conditions given by a full order on the vertices of the dual graph of the curve. Meromorphic differentials naturally arise in the description of the boundary objects, and the incidence variety compactification works just as well for the strata of meromorphic differentials, hence we take the meromorphic case into account from the beginning. In order to deal with meromorphic differentials, we consider the closure of the corresponding strata in the Hodge bundle over $\overline{\mathcal{M}}_{g,n}$ twisted by the polar part $\tilde{\mu}$ of the differentials, which we denote by $K\overline{\mathcal{M}}_g(\tilde{\mu})$ and introduce in Section 2.5.

Before we state the main result, let us first provide some motivation from several viewpoints for the reader to get a feel for the form of the answer that we get.

1.2. Motivation via complex analytic geometry. Given a pointed stable differential $(X, \omega, z_1, \dots, z_n) \in \Omega\overline{\mathcal{M}}_{g,n}$, that is a stable curve X with marked points z_1, \dots, z_n at the zeros of a stable differential ω , the question is whether it is the limit of a family of abelian differentials (X_t, ω_t) contained in a given stratum $\Omega\mathcal{M}_g(\mu)$ such that the z_i are the limits of the zeros of ω_t .

Suppose $f : \mathcal{X} \rightarrow \Delta$ is a family of abelian differentials over a disk Δ with parameter t , whose underlying curves degenerate to X at $t = 0$. If for an irreducible component X_v of X the limit

$$\omega_0 := \lim_{t \rightarrow 0} \omega_t$$

is not identically zero, then on X_v the limits of zeros of ω_t are simply the zeros of ω . Thus our goal is to extract from this family a nonzero (possibly meromorphic) differential for every irreducible component of X where ω_0 is identically zero. The analytic way to do this is to take for every X_v a suitable scaling parameter $\ell_v \in \mathbb{Z}_{\leq 0}$ such that the limit

$$\eta_v := \lim_{t \rightarrow 0} t^{\ell_v} \omega_t|_{X_v}$$

is well-defined and not identically zero. This is done in Lemma 4.1. Along this circle of ideas, we prove our main result by the plumbing techniques in Section 4.

1.3. Motivation via algebraic geometry. Now we sketch the algebro-geometric viewpoint of the above setting. Think of the family ω_t as a section of the vector bundle $f_*\omega_{\mathcal{X}^*/\Delta^*}$ of abelian differentials on the fibers over the punctured disc, where $\omega_{\mathcal{X}^*/\Delta^*}$ is the relative dualizing line bundle. The Hodge bundle $f_*\omega_{\mathcal{X}/\Delta}$ extends $f_*\omega_{\mathcal{X}^*/\Delta^*}$ to a vector bundle over the entire disc, but so does any twisting $f_*\omega_{\mathcal{X}/\Delta}(\sum c_v X_v)$ by an arbitrary integral linear combination of the irreducible components X_v of the central fiber X . Based on the idea of Eisenbud-Harris' limit linear series [EH86] (for curves of compact type), we want to choose coefficients c_v in such a way that ω_t extends over $t = 0$ to a section of the corresponding twisted dualizing line bundle, whose restriction η_v to every irreducible component X_v is not identically zero (see the discussion in [Che17] for more details). While the machinery of limit linear series for stable curves of arbitrary type is not available in full generality, our Definition 1.1 of twisted differentials works for all stable curves. It is modeled on the collection of η_v defined above.

The next question is to determine which twisted differentials $\eta = \{\eta_v\}$ on a pointed stable curve X arise as actual limits of abelian differentials (X_t, ω_t) that lie in a given stratum. First, η must have suitable zeros at the limit positions of the zeros of ω_t . Moreover if X is reducible, the limit of canonical line bundles of X_t is not unique, as it can be obtained by twisting the dualizing line bundle of X by any component X_v (treated as a divisor in the universal curve). Accordingly the limit η of ω_t on X can be regarded as a section of certain twisted dualizing line bundle, which is not identically zero on any component of X (for otherwise we can twist off such a component). Since the dualizing line bundle of X at a node is generated by differentials with simple poles, after twisting by $\sum c_v X_v$, on one branch of the node the zero or pole order gains c_v and on the other branch it loses c_v , hence the zero and pole orders of η on the two branches of every node must add up to the original sum of vanishing orders $-2 = (-1) + (-1)$. See [Che17, Section 4.1] for some examples and more details. We can then partially orient the dual graph Γ of X by orienting the edge from the zero of η to the pole, and leaving it unoriented if the differential has a simple pole at both branches. In this way, we obtain a partial order on the vertices v of Γ , with equality permitted (see also [FP16]).

It turns out that a partial order is insufficient to characterize actual limits of abelian differentials in a given stratum. For the degenerating family (X_t, ω_t) , comparing the scaling parameters ℓ_v discussed in Section 1.2 extends this partial order to a full order on the vertices of Γ , again with equality permitted. The final ingredient of our answer is the global residue condition (4) in Definition 1.2 that requires the limit twisted differentials η to be compatible with the full order on Γ . Simply speaking, this global residue condition arises from applying Stokes' formula to ω_t on each level of Γ (i.e., truncating Γ at vertices that are equal in the full order), for $t \rightarrow 0$.

1.4. Motivation via flat surfaces. Degeneration of flat surfaces has been studied in connection with counting problems, for example in [EMZ03; EMR12; EKZ14]. Most of the degeneration arguments there rely on a theorem of Rafi [Raf07] on the comparison of flat and hyperbolic lengths for surfaces near the boundary. Rafi used a thick-thin decomposition of the flat surfaces by cutting along hyperbolically short curves. For each piece of the thick-thin decomposition Rafi defined a real number, the size, in terms of hyperbolic geometry. His main theorem says that after rescaling by size, hyperbolic and flat lengths on the thick pieces are comparable, up to universal constants. Rafi's size is closely related to the scaling parameters ℓ_v discussed above, implied by comparing our scaling limit (4.1) with the 'geometric compactification theorem' [EKZ14, Theorem 10].

Rafi associated to every closed geodesic a flat representative of an annular neighborhood. Depending on the curvature of the boundary such a neighborhood is composed of flat cylinders in the middle and expanding annuli on both sides, any of the three possibly not being present. For degenerating families of flat surfaces this observation can be applied to the vanishing cycles of the family nearby a nodal fiber. In our result we use the vanishing cycles to read off the global residue condition that constrains degeneration of flat surfaces.

While some of our terminology might be translated into the language of [Raf07] or [EKZ14], in the literature on flat surfaces and Teichmüller dynamics, no systematic

attempt to describe the set of all possible limit objects under degeneration has been made. In Section 5 we provide an alternative proof of our main result by constructions of flat surfaces, where a pair of half-infinite cylinders correspond to two simple poles of a twisted differential attached together, and an expanding annulus that appears corresponds to a zero matching a higher order pole.

1.5. Level graphs. We now introduce the relevant notions that will allow us to state our result. Recall that Γ denotes the dual graph of a nodal curve X , whose vertices and edges correspond to irreducible components and nodes of X , respectively. First, we start with the idea of comparing irreducible components of X .

A *full order* on the graph Γ is a relation \succsim on the set V of vertices of Γ that is reflexive, transitive, and such that for any $v_1, v_2 \in V$ at least one of the statements $v_1 \succsim v_2$ or $v_2 \succsim v_1$ holds. We say that v_1 is of *higher or equal level* compared with v_2 if and only if $v_1 \succsim v_2$. We write $v_1 \asymp v_2$ if they are of the same level, that is if both $v_1 \succsim v_2$ and $v_2 \succsim v_1$ hold. We write $v_1 \succ v_2$ if $v_1 \succsim v_2$ but $v_2 \not\succsim v_1$, and say that v_1 is of *higher level* than v_2 . We call the set of maxima of V the *top level* vertices.

We remark that equality is permitted in our definition of a full order. Any map $\ell : V \rightarrow \mathbb{R}$ assigning real numbers to vertices of Γ defines a full order on Γ by setting $v_1 \succsim v_2$ if and only if $\ell(v_1) \geq \ell(v_2)$. Conversely, every full order can be induced from such a level map, but not from a unique one. Later we will see that the levels are related to the scaling parameters introduced in Section 1.2, hence it would be convenient to consider the maps $\ell : V \rightarrow \mathbb{R}_{\leq 0}$ assigning *non-positive* levels only, with $\ell^{-1}(0) \neq \emptyset$ being the top level.

We call a graph Γ equipped with a full order on its vertices a *level graph*, denoted by $\bar{\Gamma}$. We will use the two notions ‘full order’ and ‘level graph’ interchangeably, and draw the level graphs by aligning horizontally vertices of the same level, so that a level map is given by the projection to the vertical axis, and the top level vertices are actually placed at the top (see the examples in Section 3).

1.6. Twisted differentials. Throughout the paper we use $\text{ord}_q \eta$ to denote the zero or pole order of a differential η at q , and use $\text{Res}_q \eta$ to denote the residue of η at q . We now introduce the key notion of twisted differentials.

Definition 1.1. For a tuple of integers $\mu = (m_1, \dots, m_n)$, a *twisted differential of type μ* on a stable n -pointed curve (X, z_1, \dots, z_n) is a collection of (possibly meromorphic) differentials η_v on the irreducible components X_v of X such that no η_v is identically zero and the following properties hold:

- (0) (**Vanishing as prescribed**) Each differential η_v is holomorphic and nonzero outside of the nodes and marked points of X_v . Moreover, if a marked point z_i lies on X_v , then $\text{ord}_{z_i} \eta_v = m_i$.
- (1) (**Matching orders**) For any node of X that identifies $q_1 \in X_{v_1}$ with $q_2 \in X_{v_2}$,

$$\text{ord}_{q_1} \eta_{v_1} + \text{ord}_{q_2} \eta_{v_2} = -2.$$

- (2) (**Matching residues at simple poles**) If at a node of X that identifies $q_1 \in X_{v_1}$ with $q_2 \in X_{v_2}$ the condition $\text{ord}_{q_1} \eta_{v_1} = \text{ord}_{q_2} \eta_{v_2} = -1$ holds, then $\text{Res}_{q_1} \eta_{v_1} + \text{Res}_{q_2} \eta_{v_2} = 0$.

These conditions imply that the set of zeros and poles of a twisted differential consists of the marked points z_i and some of the nodes of the curve X .

1.7. Twisted differentials compatible with a level graph. We want to study under which conditions a twisted differential arises as a limit in a degenerating family of abelian differentials contained in a given stratum. As mentioned before, the conditions depend on a full order on the dual graph Γ of X , and we need a little more notation.

Suppose that $\bar{\Gamma}$ is a level graph with the full order determined by a level function ℓ . For a given level L we call the subgraph of Γ that consists of all vertices v with $\ell(v) > L$ along with edges between them the *graph above level L* of Γ , and denote it by $\bar{\Gamma}_{>L}$. We similarly define the graph $\bar{\Gamma}_{\geq L}$ *above or at level L* , and the graph $\bar{\Gamma}_{=L}$ *at level L* .

Accordingly we denote by $X_{>L}$ the subcurve of X with dual graph $\bar{\Gamma}_{>L}$ etc. For any node q connecting two irreducible components X_v and $X_{v'}$, if $v \succ v'$, we denote by $q^+ \in X_v$ and $q^- \in X_{v'}$ the two preimages of the node on the normalization of X . If $v \asymp v'$, we still write q_v^\pm for the preimages of the node, where the choice of which one is q_v^+ is arbitrary and will be specified. In the same way we denote by $v^+(q)$ and $v^-(q)$ the vertices at the two ends of an edge representing a node q .

Definition 1.2. Let (X, z_1, \dots, z_n) be an n -pointed stable curve with a level graph $\bar{\Gamma}$. A twisted differential η of type μ on X is called *compatible with $\bar{\Gamma}$* if, in addition to conditions (0), (1), (2) in Definition 1.1, it also satisfies the following conditions:

- (3) **(Partial order)** If a node of X identifies $q_1 \in X_{v_1}$ with $q_2 \in X_{v_2}$, then $v_1 \succ v_2$ if and only if $\text{ord}_{q_1} \eta_{v_1} \geq -1$. Moreover, $v_1 \asymp v_2$ if and only if $\text{ord}_{q_1} \eta_{v_1} = -1$.
- (4) **(Global residue condition)** For every level L and every connected component Y of $X_{>L}$ that does not contain a marked point with a prescribed pole (i.e., there is no $z_i \in Y$ with $m_i < 0$) the following condition holds. Let q_1, \dots, q_b denote the set of all nodes where Y intersects $X_{=L}$. Then

$$\sum_{j=1}^b \text{Res}_{q_j^-} \eta_{v^-(q_j)} = 0,$$

where we recall that $q_j^- \in X_{=L}$ and $v^-(q_j) \in \bar{\Gamma}_{=L}$.

We point out that a given twisted differential satisfying conditions (0), (1) and (2) may not be compatible with any level graph, or may be compatible with different level graphs with the same underlying dual graph. Condition (3) is equivalent to saying that, if v_1 is of higher level than v_2 , then η_{v_1} is holomorphic at every node of the intersections of X_{v_1} and X_{v_2} , and moreover if $v_1 \asymp v_2$, then η_{v_1} and η_{v_2} have simple poles at every node where they intersect.

1.8. Main result. Recall that the incidence variety compactification of a stratum of abelian differentials $\mathbb{P}\Omega\mathcal{M}_g(\mu)$ is defined as the closure of the stratum in the projectivized Hodge bundle over $\overline{\mathcal{M}}_{g,n}$ that parameterizes pointed stable differentials. For a stratum of meromorphic differentials, we take the closure in the twisted Hodge bundle by the polar part of the differentials (see Section 2 for details). Our main result characterizes boundary points of the incidence variety compactification for both cases.

Theorem 1.3. *A pointed stable differential $(X, \omega, z_1, \dots, z_n)$ is contained in the incidence variety compactification of a stratum $\mathbb{P}\Omega\mathcal{M}_g(\mu)$ if and only if the following conditions hold:*

- (i) *There exists a level graph $\bar{\Gamma}$ on X such that its maxima are the irreducible components X_v of X on which ω is not identically zero.*
- (ii) *There exists a twisted differential η of type μ on X , compatible with $\bar{\Gamma}$.*
- (iii) *On every irreducible component X_v where ω is not identically zero, $\eta_v = \omega|_{X_v}$.*

In Sections 1.2 and 1.3, we have briefly explained the ideas behind the necessity of these conditions. The sufficiency part is harder, that is, how can we deform pointed stable differentials satisfying the above conditions into the interior of the stratum? We provide two proofs, in Section 4 by using techniques of plumbing in complex analytic geometry, and in Section 5 by using constructions of flat surfaces.

Recall that the incidence variety compactification combines the two approaches of compactifying the strata in the projectivized Hodge bundle $\mathbb{P}\Omega\overline{\mathcal{M}}_g$ over $\overline{\mathcal{M}}_g$ and in the Deligne-Mumford space $\overline{\mathcal{M}}_{g,n}$. In particular, it admits two projections π_1 to $\mathbb{P}\Omega\overline{\mathcal{M}}_g$ and π_2 to $\overline{\mathcal{M}}_{g,n}$ by forgetting the marked points and forgetting the differentials, respectively. Hence our result completely determines the strata closures in the Hodge bundle compactification and in the Deligne-Mumford compactification.

Corollary 1.4. *A stable differential (X, ω) lies in the Hodge bundle compactification of a stratum if and only if there exists a pointed stable differential satisfying the conditions in Theorem 1.3 that maps to (X, ω) via π_1 .*

A pointed stable curve (X, z_1, \dots, z_n) lies in the Deligne-Mumford compactification of a stratum if and only if there exists a pointed stable differential satisfying the conditions in Theorem 1.3 that maps to (X, z_1, \dots, z_n) via π_2 .

Remark 1.5. Given a level graph $\bar{\Gamma}$, there can only exist finitely many twisted differentials η compatible with $\bar{\Gamma}$ and satisfying the conditions of the theorem, up to scaling η_v on each irreducible component by a nonzero number. Indeed, if all the pole and zero orders of η at all nodes are given, it determines η uniquely up to scaling on each irreducible component. For any irreducible component X_v on the bottom level, the zeros and poles of η_v are prescribed outside of the nodes of X_v , and at those nodes η_v only has poles. As the total number of zeros and poles of η_v , counted with multiplicity, is equal to $2g_v - 2$, it implies that the sum of the orders of poles of η_v at all nodes of X_v is fixed, and hence there are finitely many choices. For each such choice, on every irreducible component intersecting X_v the order of zero of η at any node where it intersects X_v is thus uniquely determined. If another component on the bottom level intersects X_v , then η has a simple pole at both branches of that node. Therefore, we can prove that there are finitely many choices of η up to scaling, by induction on the number of irreducible components of X .

For later use we relate a twisted differential η and a pointed stable differential $(X, \omega, z_1, \dots, z_n)$ satisfying Theorem 1.3 as follows.

Definition 1.6. Given a twisted differential η compatible with a level graph on X , define the *associated pointed stable differential* $(X, \omega, z_1, \dots, z_n)$ by taking ω to be equal

to η on all top level components and identically zero on components of lower levels, and by taking z_1, \dots, z_n to be the set of zeros and poles of η away from the nodes of X .

Conversely given a pointed stable differential $(X, \omega, z_1, \dots, z_n)$, if it is associated to a twisted differential η compatible with certain level graph on X , we say that η is an *associated twisted differential* of $(X, \omega, z_1, \dots, z_n)$.

Using the above definition, we can restate Theorem 1.3 as that a pointed stable differential $(X, \omega, z_1, \dots, z_n)$ is contained in the incidence variety compactification of $\mathbb{P}\Omega\mathcal{M}_g(\mu)$ if and only if there exists a level graph and a compatible twisted differential η of type μ such that $(X, \omega, z_1, \dots, z_n)$ is associated with η .

Note that multiplying η by any nonzero numbers on components not of top level does not change the associated pointed stable differential. Hence given a pointed stable differential, the associated twisted differential η may not be unique. Indeed it is not unique even up to scaling on each irreducible component (see Example 3.2).

1.9. History of the project and related work. Recently there have been several attempts via different viewpoints that aim at understanding the boundary behavior of the strata of abelian differentials. In a talk given in August 2008, Kontsevich discussed the problem of compactifying the strata, focusing on the matching order and matching residue conditions. In [Gen18] the third author studied the incidence variety compactification and applied the plumbing techniques to prove a special case of our main result when all the residues of a twisted differential are zero. In that case the global residue condition (4) obviously holds. Motivated by the theory of limit linear series, in [Che17] the second author studied the Deligne-Mumford strata compactification and deduced the necessity of the conditions (0), (1), and (2). He also obtained partial smoothing results in the case of curves of pseudo-compact type by combining techniques of algebraic geometry and flat geometry. In [FP16] Farkas and Pandharipande studied the Deligne-Mumford strata compactification by imposing the conditions (0), (1), (2), and (3), i.e., without the global residue condition (4). It turns out the corresponding loci are reducible in general, containing extra components of equal dimension or one less in the boundary of $\overline{\mathcal{M}}_{g,n}$. Modulo a conjectural relation to Pixton's formula of the double ramification cycle, in [FP16, Appendix] Janda, Pandharipande, Pixton, and Zvonkine used the extra components to recursively compute the cycle classes of the strata in $\overline{\mathcal{M}}_{g,n}$. In [MW17] Mirzakhani and Wright concentrated on a collapse of the Hodge bundle compactification by keeping track only of components where the stable differentials are not identically zero. They proved an identification between the tangent space of the boundary of a $\mathrm{GL}_2^+(\mathbb{R})$ -orbit closure and the intersection of the tangent space to the orbit closure with the tangent space to the boundary of their compactification. In [GKN17] Krichever, Norton, and the fourth author studied degenerations of meromorphic differentials with all periods real, where plumbing techniques are also used and a full order on the dual graph also arises.

In Summer 2015 the authors of the current paper met in various combinations on several occasions, including in Bonn, Luminy, Salt Lake City, and Boston. After stimulating discussions, the crucial global residue condition and the proof of sufficiency emerged, finally completing the characterization of the compactification of the strata.

1.10. Applications. The main novel aspect of our determination of the closures of the strata is the global residue condition, which is used to characterize exactly those stable differentials that appear in the closure of a stratum in the Hodge bundle, and not to extraneous components. Thus any further work aimed at understanding the structure of strata compactifications must build on our description in an essential way. In particular, the global residue condition was a cornerstone in the work of Sauvaget [Sau17] who analyzed the boundary of the strata in the Hodge bundle in order to understand the homology classes of the strata closures, and in the work of Mullane [Mul17], who used the global residue condition to analyze certain divisor closures in $\overline{\mathcal{M}}_{g,n}$, and discovered an infinite series of new extremal effective divisors. Furthermore, in [CC18] Qile Chen and the second author described algebraically the principal boundary of the strata in terms of twisted differentials, solving a problem that had been open for flat surfaces for more than a decade, while Ulirsch, Werner, and the fifth author applied in [MUW17] our compactification to solve the realizability problem for constructing the tropical Hodge bundle, posed in [LU17]. Finally in Section 3.6, as a consequence of our result we provide an efficient description for the degeneration of Weierstraß divisors on certain binary curve, recovering a main case in the work of [EM02].

1.11. Organization of the paper. In Section 2 we review basic properties about moduli spaces of curves and abelian differentials. We further compare the incidence variety compactification to the Hodge bundle compactification and the Deligne-Mumford compactification, which illustrates what extra information the incidence variety compactification gains. In Section 3 we apply Theorem 1.3 to analyze explicitly a number of examples that characterize the significance and delicacy of the global residue condition. In Section 4 we prove Theorem 1.3 by the method of plumbing. Finally in Section 5 we provide an alternative proof by constructions of flat surfaces. In particular, Sections 3, 4 and 5 are independent. Depending on the reader's background and interests, they can be read in any order.

1.12. Acknowledgments. We are grateful to Curt McMullen for useful conversations on the proof of the main theorem, to Gavril Farkas and Rahul Pandharipande for communications with us at the Arbeitstagung at Max Planck Institute in Bonn, June 2015, and to Adrian Sauvaget and Dimitri Zvonkine for discussions at the Flat Surfaces workshop in Luminy, July 2015. We thank all the organizers and participants of the Flat Surfaces workshop for their interests in this problem and motivating us to work on the proof from the viewpoint of flat geometry. The second author thanks Alex Eskin, Joe Harris, and Anton Zorich for encouraging him over the years to study Teichmüller dynamics via algebraic geometry. The fourth author thanks Igor Krichever and Chaya Norton, with whom he has been developing related ideas and techniques for real-normalized meromorphic differentials. Finally we thank the anonymous referees for carefully reading the paper and many useful comments.

2. COMPARISON OF STRATA COMPACTIFICATIONS

In this section we introduce the basic terminology about moduli spaces of curves and strata of abelian differentials. We also define the incidence variety compactification and compare it to the Hodge bundle and the Deligne-Mumford compactifications.

2.1. Moduli spaces of curves. Denote by \mathcal{M}_g the *moduli space of curves of genus g* that parameterizes smooth and connected complex curves of genus g , up to biholomorphism. Denote by $\mathcal{M}_{g,n}$ the *moduli space of n -pointed genus g curves* that parameterizes smooth and connected complex curves of genus g together with n distinct (ordered) marked points. The space $\mathcal{M}_{g,n}$ is a complex orbifold of dimension $3g - 3 + n$.

Recall that a *stable n -pointed curve* is a connected curve with at worst nodal singularities, with n distinct marked smooth points, such that the automorphism group of the curve preserving the marked points is finite. Denote by $\overline{\mathcal{M}}_{g,n}$ the *Deligne-Mumford compactification* of $\overline{\mathcal{M}}_{g,n}$ parameterizing stable n -pointed genus g curves.

Let \mathfrak{S} be a subgroup of the symmetric group \mathfrak{S}_n . Then \mathfrak{S} acts on $\overline{\mathcal{M}}_{g,n}$ by permuting the marked points. The quotient of $\overline{\mathcal{M}}_{g,n}$ by \mathfrak{S} is denoted by $\overline{\mathcal{M}}_{g,\{n\}}^{\mathfrak{S}}$ or simply $\overline{\mathcal{M}}_{g,\{n\}}$ when the group \mathfrak{S} is clear from the context.

The *dual graph* of a nodal curve X is the graph Γ whose vertices correspond to the irreducible components of X . For every node of X joining two components v_1 and v_2 (possibly being the same component) the dual graph has an edge connecting v_1 and v_2 . For every marked point there is a leg or equivalently a half-edge attached to the vertex corresponding to the irreducible component that contains the marked point.

For a connected nodal curve X , if removing a node disconnects X , we say that it is a *separating* node. Otherwise we call it a *non-separating* node. If the two branches of a node belong to the same irreducible component of X , we say that it is an *internal* node.

2.2. Moduli spaces of abelian differentials. The *moduli space of abelian differentials* $\Omega\mathcal{M}_g^*$ is the complement of the zero section in the *Hodge bundle* $\Omega\mathcal{M}_g \rightarrow \mathcal{M}_g$, which parameterizes pairs (X, ω) where X is a smooth and connected curve of genus g , and ω is a holomorphic differential on X . The space $\Omega\mathcal{M}_g^*$ has a natural stratification according to the orders of zeros of ω . Let $\mu = (m_1, \dots, m_n)$ be a partition of $2g - 2$ by positive integers. The *stratum* $\Omega\mathcal{M}_g(\mu)$ of *abelian differentials of type μ* , as a subspace of $\Omega\mathcal{M}_g^*$, parameterizes abelian differentials (X, ω) such that ω has n distinct zeros of orders m_1, \dots, m_n , respectively.

A *stable differential* on a nodal curve X is a (possibly meromorphic) differential ω on X which is holomorphic outside of the nodes of X and which has at worst simple poles at the nodes, with opposite residues. The Hodge bundle extends to a vector bundle $\Omega\overline{\mathcal{M}}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$, the total space of the relative dualizing sheaf of the universal family $f : \mathcal{X} \rightarrow \overline{\mathcal{M}}_{g,n}$. The fiber of this vector bundle over a pointed nodal curve (X, z_1, \dots, z_n) parameterizes *pointed stable differentials* $(X, \omega, z_1, \dots, z_n)$, where ω is a stable differential on (X, z_1, \dots, z_n) .

There is a natural \mathbb{C}^* -action on the Hodge bundle by scaling the differentials. This action preserves the stratification of $\Omega\mathcal{M}_g^*$. The quotient of $\Omega\mathcal{M}_g^*$ under this action is

denoted by $\mathbb{P}\Omega\mathcal{M}_g$. In general, quotient spaces by such a \mathbb{C}^* -action will be denoted by adding the letter \mathbb{P} .

2.3. The incidence variety compactification. When abelian differentials degenerate, we want to keep track of the information about both the limit stable differentials and the limit positions of the marked zeros. This motivates the definition of the incidence variety.

For a partition $\mu = (m_1, \dots, m_n)$ of $2g-2$, the (*ordered*) *incidence variety* $\mathbb{P}\Omega\mathcal{M}_{g,n}^{\text{inc}}(\mu)$ is defined to be

$$(2.1) \quad \mathbb{P}\Omega\mathcal{M}_{g,n}^{\text{inc}}(\mu) := \left\{ (X, \omega, z_1, \dots, z_n) \in \mathbb{P}\Omega\mathcal{M}_{g,n} : \text{div}(\omega) = \sum_{i=1}^n m_i z_i \right\}.$$

The (*ordered*) *incidence variety compactification* $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,n}^{\text{inc}}(\mu)$ is defined to be the closure of the incidence variety inside $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,n}$.

If $m_i = m_j$ for some $i \neq j$, then interchanging z_i and z_j preserves the incidence variety. Hence the subgroup \mathfrak{S} of the symmetric group \mathfrak{S}_n generated by such transpositions acts on $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,n}^{\text{inc}}(\mu)$ by permuting z_1, \dots, z_n accordingly. We define the (*unordered*) *incidence variety* to be the quotient

$$(2.2) \quad \mathbb{P}\Omega\mathcal{M}_{g,\{n\}}^{\text{inc}}(\mu) := \mathbb{P}\Omega\mathcal{M}_{g,n}^{\text{inc}}(\mu)/\mathfrak{S}$$

and define the (*unordered*) *incidence variety compactification* to be the quotient

$$(2.3) \quad \mathbb{P}\Omega\overline{\mathcal{M}}_{g,\{n\}}^{\text{inc}}(\mu) := \mathbb{P}\Omega\overline{\mathcal{M}}_{g,n}^{\text{inc}}(\mu)/\mathfrak{S}.$$

Since the ordered and unordered incidence variety compactifications differ only by permuting the marked points, we call both of them the incidence variety compactification. In case we need to distinguish them, we do so by specifying n or $\{n\}$ in the subscripts.

On a smooth curve X , a non-zero differential ω determines its zeros z_i along with their multiplicities. Hence $\mathbb{P}\Omega\mathcal{M}_{g,\{n\}}^{\text{inc}}(\mu)$ is isomorphic to the projectivized stratum of abelian differentials $\mathbb{P}\Omega\mathcal{M}_g(\mu)$. In order to understand degeneration of abelian differentials in $\mathbb{P}\Omega\mathcal{M}_g(\mu)$, we need to describe the boundary points of the incidence variety compactification.

2.4. Moduli spaces of pointed meromorphic differentials. The moduli spaces of meromorphic differentials from the viewpoint of flat geometry have been investigated by Boissy [Boi15a], including their dimensions and connected components. Let

$$\mu = (m_1, \dots, m_r; m_{r+1}, \dots, m_{r+s}; m_{r+s+1}, \dots, m_{r+s+l})$$

be an n -tuple of integers such that $\sum_{i=1}^n m_i = 2g-2$, where $m_i > 0$ for $i \leq r$, $m_{r+1} = \dots = m_{r+s} = 0$, and $m_i < 0$ for $i > r+s$. We call such μ a meromorphic type, and denote by $\Omega\mathcal{M}_g(\mu)$ the *moduli space of meromorphic differentials of type μ* . It parameterizes n -pointed meromorphic differentials $(X, \omega, z_1, \dots, z_n)$ on a smooth curve X such that the order of ω at z_i is equal to m_i , which may be a zero, regular point, or pole, corresponding to whether $m_i > 0$, $m_i = 0$, or $m_i < 0$, respectively. As there are infinitely many meromorphic types for a fixed genus, these moduli spaces no

longer form a stratification of a fixed ambient space, but by a slight abuse of language we still call them *strata of meromorphic differentials*.

2.5. The incidence variety compactification in the meromorphic case. To mimic the definition in the abelian case we need to generalize the notion of the Hodge bundle. We denote the polar part of μ by $\tilde{\mu} = (m_{r+s+1}, \dots, m_n)$. We then define the *pointed Hodge bundle twisted by $\tilde{\mu}$* to be the bundle

$$K\overline{\mathcal{M}}_{g,n}(\tilde{\mu}) = f_*\omega_{\mathcal{X}/\overline{\mathcal{M}}_{g,n}}\left(-\sum_{i=r+s+1}^n m_i \mathcal{Z}_i\right)$$

over $\overline{\mathcal{M}}_{g,n}$, where we have denoted by \mathcal{Z}_i the image of the section of the universal family f given by the i 'th marked point. We call points $(X, \omega, z_1, \dots, z_n) \in K\overline{\mathcal{M}}_{g,n}(\tilde{\mu})$ *pointed stable differentials (of type $\tilde{\mu}$)*. Note that if μ is a holomorphic type, then the tuple $\tilde{\mu}$ is empty, so $K\overline{\mathcal{M}}_g(\tilde{\mu}) = \Omega\overline{\mathcal{M}}_g$, and hence it recovers the preceding setting for the abelian case.

For any meromorphic type μ , we perform the same operations inside the space $K\overline{\mathcal{M}}_g(\tilde{\mu})$ as in the abelian case. The *(ordered) incidence variety* $\mathbb{P}\Omega\mathcal{M}_{g,n}^{\text{inc}}(m_1, \dots, m_n)$ is defined as

$$(2.4) \quad \mathbb{P}\Omega\mathcal{M}_{g,n}^{\text{inc}}(\mu) = \left\{ (X, \omega, z_1, \dots, z_n) \in \mathbb{P}K\overline{\mathcal{M}}_{g,n}(\tilde{\mu}) : \text{div}(\omega) = \sum_{i=1}^n m_i z_i \right\},$$

and the *(ordered) incidence variety compactification* $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,n}^{\text{inc}}(\mu)$ is defined to be its closure in $\mathbb{P}K\overline{\mathcal{M}}_{g,n}(\tilde{\mu})$. We define the *(unordered) incidence variety* to be the quotient

$$(2.5) \quad \mathbb{P}\Omega\mathcal{M}_{g,\{n\}}^{\text{inc}}(\mu) = \mathbb{P}\Omega\mathcal{M}_{g,n}^{\text{inc}}(\mu)/\mathfrak{S},$$

where \mathfrak{S} is defined as in Section 2.3. Finally, we define the *(unordered) incidence variety compactification* to be the quotient

$$(2.6) \quad \mathbb{P}\Omega\overline{\mathcal{M}}_{g,\{n\}}^{\text{inc}}(\mu) = \mathbb{P}\Omega\overline{\mathcal{M}}_{g,n}^{\text{inc}}(\mu)/\mathfrak{S}.$$

Again, by a slight abuse of language we sometimes skip the terms ‘ordered’ and ‘unordered’, and refer to both of them as the incidence variety compactification of the strata of meromorphic differentials.

2.6. Comparison to the Hodge bundle and the Deligne-Mumford compactifications. As said earlier, the incidence variety compactification admits two forgetful maps to the projectivized Hodge bundle and to the Deligne-Mumford space, respectively. In this subsection, we discuss these maps and demonstrate that they indeed forget information. For simplicity of notation, we will only state this in the holomorphic case. The discussion can be easily generalized to the meromorphic case. Thus we start with a partition $\mu = (m_1, \dots, m_n)$ of $2g - 2$ by positive integers.

The forgetful map

$$(2.7) \quad \pi_1 : \mathbb{P}\Omega\overline{\mathcal{M}}_{g,\{n\}}^{\text{inc}}(\mu) \rightarrow \mathbb{P}\Omega\overline{\mathcal{M}}_g(\mu)$$

forgets the marked points z_1, \dots, z_n . More precisely, the image of $(X, \omega, z_1, \dots, z_n)$ under π_1 is (X', ω') , where X' is obtained from X (as an unmarked curve) by blowing

down all \mathbb{P}^1 tails and bridges, and ω' can be identified with the restriction of ω to the remaining components (see [Gen18, Lemma 2.4] for details). The other forgetful map

$$(2.8) \quad \pi_2 : \mathbb{P}\Omega\overline{\mathcal{M}}_{g,\{n\}}^{\text{inc}}(\mu) \rightarrow \overline{\mathcal{M}}_{g,\{n\}}$$

forgets the stable differential ω , hence the image of $(X, \omega, z_1, \dots, z_n)$ under π_2 is just (X, z_1, \dots, z_n) .

Since a differential on a compact curve is determined uniquely (up to scaling) by the locations and orders of its zeros, both maps π_1 and π_2 , when restricted to $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,\{n\}}^{\text{inc}}(\mu)$, are isomorphisms onto their respective images. However, over the boundary the fibers of both maps can be more complicated. In particular, they may no longer be finite, and neither image dominates the other.

Proposition 2.1. *For $g \geq 3$ and $n \geq 2$, the following properties hold:*

- (i) *The map π_1 is not finite, and there does not exist a map*

$$f : \mathbb{P}\Omega\overline{\mathcal{M}}_g(\mu) \rightarrow \overline{\mathcal{M}}_{g,\{n\}}$$

such that $f \circ \pi_1 = \pi_2$.

- (ii) *The map π_2 is not finite, and there does not exist a map*

$$h : \pi_2\left(\mathbb{P}\Omega\overline{\mathcal{M}}_{g,\{n\}}^{\text{inc}}(\mu)\right) \rightarrow \mathbb{P}\Omega\overline{\mathcal{M}}_g(\mu)$$

such that $h \circ \pi_2 = \pi_1$.

This proposition is not a priori clear. Indeed it uses the full strength of Theorem 1.3 about characterizing the boundary points of $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,\{n\}}^{\text{inc}}(\mu)$. As a result we see that the incidence variety compactification contains more information than both the Hodge bundle compactification and the Deligne-Mumford compactification of the strata.

Proof. We first prove (i). Suppose X is the union of an elliptic curve X_1 with a curve X_{g-1} of genus $g-1$ intersecting at a node $q^+ \sim q^-$ such that $(2g-4)q^+ \sim K_{X_{g-1}}$ and that X_1 contains all the marked points z_1, \dots, z_n . Put X_{g-1} on a higher level than X_1 . The corresponding level graph $\overline{\Gamma}$ of X is represented on the left side of Figure 1. Take a stable differential ω on X such that $\omega|_{X_{g-1}}$ is a holomorphic differential with a unique zero at q^+ of multiplicity $2g-4$, and $\omega|_{X_1}$ is identically zero. Take a twisted differential η on X such that $\eta_{X_{g-1}} = \omega|_{X_{g-1}}$ and such that η_{X_1} is a meromorphic differential with $\text{div}(\eta_{X_1}) = \sum_{i=1}^n m_i z_i - (2g-2)q^-$.

One checks that η satisfies all the conditions in Definitions 1.1 and 1.2. In particular, the global residue condition follows from the residue theorem on X_1 , because η_{X_1} has a unique pole at q^- , and hence $\text{Res}_{q^-} \eta_{X_1} = 0$. Since η is compatible with $\overline{\Gamma}$ and $(X, \omega, z_1, \dots, z_n)$ is the associated pointed stable differential of η (see Definition 1.6), by Theorem 1.3 $(X, \omega, z_1, \dots, z_n)$ is contained in the incidence variety compactification $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,\{n\}}^{\text{inc}}(\mu)$. It implies that the π_1 -preimage of the stable differential (X, ω) is isomorphic to

$$\left\{ (z_1, \dots, z_n) \in (X_1)^n \setminus \Delta : \sum_{i=1}^n m_i z_i = (2g-2)q^- \right\} / \mathfrak{S},$$

where Δ is the big diagonal parameterizing the tuples where at least two marked points coincide. This preimage has dimension $n-1$. Since the π_2 -image retains the information about the positions of the marked points, this example implies that there does not exist a map f such that $f \circ \pi_1 = \pi_2$.




FIGURE 1. The level graphs used in the proof of Proposition 2.1

Next we prove (ii). Let X be the union of two elliptic curves X_1 , X_2 and a curve X_{g-2} of genus $g-2$, whose dual graph with a chosen full order $\bar{\Gamma}$ is represented on the right side of Figure 1. In this level graph X_1 and X_{g-2} are on the top level, both higher than X_2 . Further suppose that $(2g-6)q_2^+ \sim K_{X_{g-2}}$ and that the points $z_1, \dots, z_n \in X_2$ are chosen such that there exists a meromorphic differential η_{X_2} on X_2 with $\text{div}(\eta_{X_2}) = \sum_{i=1}^n m_i z_i - 2q_1^- - (2g-4)q_2^-$ and such that $\text{Res}_{q_1^-} \eta_{X_2} = \text{Res}_{q_2^-} \eta_{X_2} = 0$. The existence of such η_{X_2} is proved in [Boi15b]. Then (X, z_1, \dots, z_n) specifies a point in $\overline{\mathcal{M}}_{g, \{n\}}$. Take a twisted differential η on X such that η_{X_1} on X_1 is holomorphic and nowhere vanishing, η_{X_2} is given as above on X_2 , and $\eta_{X_{2g-2}}$ on X_{2g-2} satisfies $\text{div}(\eta_{X_{2g-2}}) = (2g-6)q_2^+$.

One checks that η satisfies all the required conditions to be compatible with $\bar{\Gamma}$. In particular, the global residue condition holds because η_{X_2} has zero residues at both q_1^- and q_2^- . Hence by Theorem 1.3 the associated pointed stable differential $(X, \omega, z_1, \dots, z_n)$ is contained in the incidence variety compactification $\mathbb{P}\Omega\overline{\mathcal{M}}_{g, \{n\}}^{\text{inc}}(\mu)$, where $\omega|_{X_1} = \eta_{X_1}$, $\omega|_{X_{g-2}} = \eta_{X_{g-2}}$, and ω is identically zero on X_2 . Note that one can scale η on the top level components X_1 and X_{g-2} by a pair of nonzero scalars $(\lambda_1, \lambda_{g-2})$, which does not affect its compatibility with $\bar{\Gamma}$. The associated stable differential ω is scaled accordingly on X_1 and X_{g-2} , but the underlying marked curve (X, z_1, \dots, z_n) remains the same. In other words, the π_2 -preimage of (X, z_1, \dots, z_n) in the incidence variety compactification contains the space of the projectivized pairs $[\lambda_1, \lambda_{g-2}]$, which is one-dimensional. Since the image of $(X, \omega, z_1, \dots, z_n)$ under π_1 retains the scaling information, there does not exist a map h such that $h \circ \pi_2 = \pi_1$ holds. \square

Remark 2.2. Conceptually speaking, the map π_2 fails to be injective for two reasons. First, for a component of X , there may exist a nontrivial linear equivalence relation between the marked points in that component. For example, let X be the union of a hyperelliptic curve Y of genus $g-1$, and a \mathbb{P}^1 component, intersecting at two points q_1 and q_2 which are Weierstraß points of Y . Moreover suppose that all the marked points are contained in \mathbb{P}^1 . Then (X, z_1, \dots, z_n) can be the image of $(X, \omega, z_1, \dots, z_n)$ under π_2 , where ω is identically zero on \mathbb{P}^1 and restricts to Y as a differential with a zero of order $2k$ at q_1 and a zero of order $2(g-k-2)$ at q_2 , for any $k \in \{0, \dots, g-2\}$. The other

reason is that some scaling factors for the differentials on the top level components are lost, as discussed in the second part of the proof of Proposition 2.1 (also see Lemma 2.3 below).

When we analyze the examples presented in Section 3, the information of the dimension of fibers of π_2 will play a significant role. Hence we conclude this section by the following observation.

Lemma 2.3. *Let Δ° be an open boundary stratum of $\overline{\mathcal{M}}_{g,n}$ parameterizing nodal curves with a given dual graph. Let (X, z_1, \dots, z_n) be a curve in the intersection of the locus $\pi_2(\mathbb{P}\Omega\overline{\mathcal{M}}_{g,n}^{\text{inc}}(\mu))$ with Δ° . Then the dimension of the fiber of π_2 over (X, z_1, \dots, z_n) is one less than the maximal number of connected components of the graph $\Gamma_{=0}$, where the maximum is taken over all level graph structures $\overline{\Gamma}$ on Γ , such that there exists a compatible twisted differential of type μ on (X, z_1, \dots, z_n) .*

Proof. Take a twisted differential η on X compatible with a chosen level graph $\overline{\Gamma}$. Suppose $(X, \omega, z_1, \dots, z_n)$ is the pointed stable differential associated to $(\eta, \overline{\Gamma})$. Then ω is equal to η restricted to all top level components, and is identically zero elsewhere. The image of $(X, \omega, z_1, \dots, z_n)$ under π_2 then only retains the information on the zeroes of ω on all the irreducible components of top level, and thus loses the information of the individual scaling factors of η on each such irreducible component. However, if two such top level irreducible components are connected by an edge, the matching residue condition at the corresponding node prescribes that the scale of η on these two components is equal, and thus there is only one scale parameter lost for each connected component of the graph of the top level components. The space of such scaling factors has projective dimension equal to the number of top level components minus one. The desired conclusion thus follows from applying this analysis to all possible level graphs and compatible twisted differentials. \square

3. EXAMPLES OF THE INCIDENCE VARIETY COMPACTIFICATION

To illustrate all the aspects of the incidence variety compactification, in this section we study many examples. We start with examples discussing possible choices of level graphs and compatible twisted differentials for a fixed dual graph, thus showing that there are indeed choices involved, and that all our data are necessary. We then describe in detail the incidence variety compactification for a number of strata in low genus.

Throughout this section, we use the following notation. Denote by K_{X_i} the canonical line bundle of an irreducible component X_i of a nodal curve X . A node joining two irreducible components X_i and X_j is denoted by q_k . Moreover, if $X_i \succ X_j$, then the node q_k is obtained by identifying the points $q_k^+ \in X_i$ with $q_k^- \in X_j$. We denote by Γ the dual graph of X , and by $\overline{\Gamma}$ a full order on Γ . For a twisted differential η and a stable differential ω on X , we use η_i and ω_i to denote their restrictions to the component X_i , respectively. We also remind the reader to review Definition 1.6 for η and ω being associated with each other.

3.1. Cautionary examples. We present some examples that serve as an illustration for our formulation of the conditions on twisted differentials as well as a warning regarding the extent to which the choices of $\overline{\Gamma}$ and η determine each other.

Example 3.1. (*Twisted differentials do not automatically satisfy the global residue condition*) Let X be a curve with three components $X_1 \asymp X_2 \succ X_3$ as represented in Figure 2. Suppose that X_1 and X_2 contain no marked poles, and suppose that $g_3 = 0$. Suppose that the only marked point on X_3 is a marked zero z_1 , i.e. $m_1 > 0$.




FIGURE 2. The curve and level graph used in Example 3.1

Let η be a twisted differential on X compatible with this level graph. Since X_3 is on the bottom level, η_3 has poles at q_1^- and q_2^- . Let r_1 and r_2 be the residues of η_3 at q_1^- and q_2^- , respectively. The residue theorem on $X_3 \cong \mathbb{P}^1$ says that $r_1 + r_2 = 0$, with no further constraints. However, the global residue condition applied to the level of X_3 implies that $r_1 = 0$ and $r_2 = 0$, which does not follow from the relation $r_1 + r_2 = 0$.

Example 3.2. (*Non-uniqueness of associated twisted differentials*) Theorem 1.3 says that a pointed stable differential lies in the incidence variety compactification of a given stratum if and only if there exists an associated twisted differential of the given type compatible with certain level graph (see Definition 1.6). However, such a twisted differential may not be unique (modulo scaling), even for a fixed level graph. For example, suppose X has three irreducible components $X_1 \succ X_2 \succ X_3$, where X_1 intersects X_2 at one point q_1 , and X_2 and X_3 intersect at two points q_2 and q_3 , and suppose all marked points z_1, \dots, z_n lie on X_3 , see Figure 3.




FIGURE 3. The curve and level graph used in Example 3.2

Suppose η is a twisted differential compatible with the corresponding level graph $\bar{\Gamma}$. Because X_1 has no marked points and it is on the top level, η_1 on X_1 is holomorphic, and it has a unique zero at q_1^+ whose order is $2g_1 - 2$. Moreover, η_3 on X_3 has all the prescribed zeros or poles at z_i , hence the sum of its pole orders at q_2^- and q_3^- is equal to $2g - 2g_3$. Suppose that η_3 has a pole of order k at q_2^- and a pole of order $2g - 2g_3 - k$ at q_3^- . Then η_2 as a differential on X_2 has a pole of order $2g_1$ at q_1^- , and has zeros of orders $k - 2$ and $2g - 2g_3 - k - 2$ at q_2^+ and q_3^+ , respectively. Finally applying the global residue condition to each level of $\bar{\Gamma}$, respectively, it says that the sum of residues of η_i on each irreducible component X_i is zero, which follows from the residue theorem. Hence in this case the global residue condition imposes no further constraints on η .

By the above analysis, there exists a twisted differential η compatible with $\bar{\Gamma}$ if and only if the pointed curve (X, z_1, \dots, z_n) satisfies

$$\begin{aligned} K_{X_1} &\sim (2g_1 - 2) q_1^+, \\ K_{X_2} &\sim (k - 2)q_2^+ + (2g - 2g_3 - k - 2)q_3^+ - 2g_1q_1^-, \\ K_{X_3} &\sim \sum_{i=1}^n m_i z_i - kq_2^- - (2g - 2g_3 - k)q_3^-. \end{aligned}$$

For special curves X these conditions can be satisfied for different values of k . For instance, take all X_i to be hyperelliptic curves of high genus, all q_i^\pm to be Weierstraß points, and k to be even. In that case we obtain a number of distinct twisted differentials η , all of which are the same on X_1 (modulo scaling) but are different on X_2 and X_3 (even after modulo scaling).

Suppose ω is the associated stable differential of η . Then $\omega_1 = \eta_1$ is determined on the top level component X_1 , which is the same for all the different choices of η (modulo scaling). Nevertheless, ω_2 and ω_3 are identically zero on the lower level components X_2 and X_3 , respectively. Hence in this case different values of k in the above give rise to distinct twisted differentials η , but the associated stable differential $(X, \omega, z_1, \dots, z_n)$ remains to be the same.

Example 3.3. (*Pointed stable differentials do not determine the level graph*) The previous example shows that there may be many different twisted differentials that are associated with a given pointed stable differential in the incidence variety compactification. We now show that a pointed stable differential $(X, \omega, z_1, \dots, z_n)$ does not necessarily determine a full order on the dual graph. The reason is that ω is identically zero on the lower level components of X , hence the lower level components may be ordered differently. An example illustrated in Figure 4 is given by a triangular subgraph on lower levels, attached to a top level component X_1 . The only two marked points are $z_1 \in X_3$ and $z_2 \in X_4$. The two different level graphs are obtained by switching the ordering of X_3 and X_4 on the bottom two levels.




FIGURE 4. The curve with two different level graphs used in Example 3.3

Suppose the genera of X_1 and X_2 are sufficiently high so that $m_1 > 2g_3 - 2$ and $m_2 > 2g_4 - 2$. Take a twisted differential η such that

$$\begin{aligned}\operatorname{div}(\eta_1) &= (2g_1 - 2)q_1^+, \\ \operatorname{div}(\eta_2) &= kq_2^+ + (2g_2 + 2g_1 - k - 2)q_3^+ - 2g_1q_1^-, \\ \operatorname{div}(\eta_3) &= m_1z_1 + (2g_3 + k - m_1)q_4 - (k + 2)q_2^-, \\ \operatorname{div}(\eta_4) &= m_2z_2 + (k - 2g_1 - 2g_2)q_3^- + (m_1 - 2 - k - 2g_3)q_4,\end{aligned}$$

where $0 \leq k \leq 2g_2 + 2g_1 - 2$. The existence of such η is equivalent to the linear equivalence conditions $\operatorname{div}(\eta_i) \sim K_{X_i}$ for all i . One can take X_i to be hyperelliptic curves, z_i and q_j to be Weierstraß points, m_i and k to be even, so that these conditions hold. In that case, if we choose suitable k such that $2g_3 + k - m_1 > -1$, then η_3 is holomorphic at q_4 on X_3 , hence η is compatible with the level graph on the right side of Figure 4, where the global residue condition follows from the residue theorem on each lower level component. Conversely if $2g_3 + k - m_1 < -1$, then η_3 has a higher order pole at q_4 on X_3 , hence η is compatible with the level graph in the middle of Figure 4. Nevertheless, in both cases the associated stable differentials ω are the same (modulo scaling on the top level component X_1), because ω is identically zero on the components X_2 , X_3 , and X_4 .

Example 3.4. (*Different level graphs give different global residue conditions*) Consider two level graphs $\bar{\Gamma}_1$ and $\bar{\Gamma}_2$ with the same underlying dual graph Γ of a curve X , presented in Figure 5. The orientation of the edges is going downwards. Because X_5 and X_6 are disjoint, $\bar{\Gamma}_1$ and $\bar{\Gamma}_2$ determine the same partial order on Γ . Further suppose there is no marked pole in the smooth locus of X .




FIGURE 5. Two level graphs $\bar{\Gamma}_1$ and $\bar{\Gamma}_2$ used in Example 3.4

We will now determine the conditions necessary for a twisted differential η to be compatible with $\bar{\Gamma}_1$ or $\bar{\Gamma}_2$, respectively. For a node q_k joining X_i and X_j with $X_i \succ X_j$, we denote by $r_{i,j}$ the residue of η_j at q_k^- . In both cases η_1 , η_2 , and η_3 are holomorphic, and η_4 , η_5 , and η_6 are meromorphic. Applying the residue theorem to η_i on X_i for $i = 4, 5, 6$, we obtain that

$$r_{1,4} + r_{3,4} = 0, \quad r_{1,5} + r_{2,5} + r_{3,5} = 0, \quad r_{2,6} + r_{4,6} = 0.$$

The global residue condition can be imposed at the bottom level and at the middle level of $\bar{\Gamma}_i$. For $\bar{\Gamma}_1$ the bottom level global residue condition implies that

$$r_{2,6} + r_{4,6} = 0$$

(since the graph above the bottom level is connected), which follows from the residue theorem. The global residue condition applied to the middle level of $\bar{\Gamma}_1$ implies that

$$r_{1,4} + r_{1,5} = 0, \quad r_{2,5} = 0, \quad r_{3,4} + r_{3,5} = 0$$

where the third condition follows from the first two and the residue theorem. Thus for $\bar{\Gamma}_1$ the global residue condition gives two extra relations among the residues of η , in addition to the residue theorem. In particular, it implies that $r_{2,5} = 0$.

For $\bar{\Gamma}_2$ the global residue condition applied to the bottom level implies that

$$r_{4,6} + r_{1,5} + r_{3,5} = 0, \quad r_{2,5} + r_{2,6} = 0$$

(since the graph above the bottom level has two connected components, one being X_2 , and the other everything else), where the two conditions are equivalent by the residue theorem. The global residue condition applied to the middle level of $\bar{\Gamma}_2$ implies that $r_{1,4} = 0$ (or equivalently, $r_{3,4} = 0$, by the residue theorem). Thus for $\bar{\Gamma}_2$ the global residue condition gives two extra relations among the residues of η , in addition to the residue theorem. In particular, it implies that $r_{1,4} = 0$, which is different from the conditions imposed by $\bar{\Gamma}_1$ in the preceding paragraph.

Example 3.5. (*The space of residues cut out by the global residue condition*) In this example we illustrate how the scaling factors of a twisted differential on lower level components of X come into play in the global residue condition. Consider the curve X with the level graph $\bar{\Gamma}$ in Figure 6. Suppose there is no marked pole in the smooth locus of X .




FIGURE 6. The curve and level graph used in Example 3.5

Let η be a twisted differential on X whose associated stable differential is ω . Suppose further that η satisfies conditions (0)–(3) with respect to $\bar{\Gamma}$ (see Definitions 1.1 and 1.2). Take a pair of scalars $\lambda = (\lambda_4, \lambda_5)$. Define a twisted differential

$$\eta(\lambda) = \{\eta_1, \eta_2, \eta_3, \lambda_4\eta_4, \lambda_5\eta_5\}.$$

Because X_4 and X_5 are the lower level components, the associated stable differential of $\eta(\lambda)$ is ω , and $\eta(\lambda)$ also satisfies conditions (0)–(3) with respect to $\bar{\Gamma}$. Conversely, any twisted differential with associated stable differential equal to ω is of the type $\eta(\lambda)$. By Theorem 1.3, $(X, \omega, z_1, \dots, z_n)$ is contained in the incidence variety compactification if and only if there exists one such $\eta(\lambda)$ satisfying the global residue condition.

Use the notation $r_{i,j}$ for residues of η as in the preceding example. Then the residues of $\eta(\lambda)$, compared to η , are multiplied by λ_4 and λ_5 on X_4 and X_5 , respectively. Applying the residue theorem to X_4 and X_5 gives

$$r_{1,4} + r_{2,4} + r_{3,4} = 0, \quad r_{1,5} + r_{2,5} + r_{3,5} = 0.$$

The global residue condition applied to $\eta(\lambda)$ and $\bar{\Gamma}$ imposes the conditions

$$\lambda_4 r_{1,4} + \lambda_5 r_{1,5} = 0, \quad \lambda_4 r_{2,4} + \lambda_5 r_{2,5} = 0, \quad \lambda_4 r_{3,4} + \lambda_5 r_{3,5} = 0.$$

By the residue theorem we can express $r_{3,4}$ and $r_{3,5}$ in terms of the other residues, hence the imposed conditions reduce to

$$\begin{cases} \lambda_4 r_{1,4} + \lambda_5 r_{1,5} = 0, \\ \lambda_4 r_{2,4} + \lambda_5 r_{2,5} = 0. \end{cases}$$

Eliminating λ_4 and λ_5 in the above further reduces the conditions to

$$r_{1,4} r_{2,5} - r_{2,4} r_{1,5} = 0.$$

Hence there exists a twisted differential $\eta(\lambda)$ compatible with $\bar{\Gamma}$ only if the residues of η satisfy this *quadratic* equation. This phenomenon will be addressed in full generality in our forthcoming work [BCGGM], where we study in detail the boundary structure of the incidence variety compactification.

3.2. Some preliminary results. We first remark that the incidence variety compactification $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,\{n\}}^{\text{inc}}(m_1, \dots, m_n)$ of any stratum has divisorial boundary (i.e., the boundary is of complex codimension one in the compactification). This follows from the fact that $\overline{\mathcal{M}}_{g,n}$ and the projectivized Hodge bundle have divisorial boundary, and this is where we take the closure of the stratum. In this section we describe explicitly the boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{g,\{n\}}^{\text{inc}}(m_1, \dots, m_n)$ for a few cases. The general method is as follows. First, we enumerate the boundary strata in $\overline{\mathcal{M}}_{g,n}$ parameterizing stable curves with a given dual graph. For each such stratum, we investigate all possible level graphs. Then we write down the space of all possible twisted differentials η that are compatible with such a level graph, which we write as a formula saying that η lies in a suitable product of strata of differentials, recording its zeros and poles on the irreducible components of X . Given such η , the associated pointed stable differential ω coincides with η on the top level components, and is identically zero elsewhere. To conclude, we use Lemma 2.3 to compute the dimension of the fiber of the map π_2 from the incidence variety compactification to its image in the Deligne-Mumford compactification (see the setting (2.8)).

For later use we denote by Δ_i° the *open* divisorial components of the boundary of $\overline{\mathcal{M}}_g$. More precisely, Δ_i° for $1 \leq i \leq [g/2]$ parameterizes reducible curves consisting of a genus i smooth component and a genus $g - i$ smooth component attached at one node, and Δ_0° parameterizes irreducible nodal curves of geometric genus $g - 1$.

We first consider the strata in genus zero, since they play a central role in many cases. Recall that there exists a meromorphic differential on \mathbb{P}^1 with any prescribed collection of zeros and poles, as long as the sum of their orders is equal to -2 . As a consequence of Theorem 1.3, we deduce the following statement about residues of meromorphic differentials on \mathbb{P}^1 .

Lemma 3.6. *Let $\Omega\mathcal{M}_0(m_0, \dots, m_n)$ be a stratum of meromorphic differentials in genus zero such that $m_0 > 0$ and $m_i < 0$ for $1 \leq i \leq n$. If $n \geq 2$, then every differential in this stratum has a nonzero residue at some pole.*

Proof. If some $m_i = -1$, then the corresponding pole is simple, hence it has a nonzero residue. From now on assume that $m_i \leq -2$ for all $1 \leq i \leq n$.

We argue by contradiction. Suppose this stratum contains a differential η_0 on $X_0 \cong \mathbb{P}^1$ with zero residue at every pole. Denote by z_0 the zero of η_0 and z_i the pole of η_0 of order m_i for $1 \leq i \leq n$. For every $i \geq 1$ we attach to X_0 at z_i a curve X_i as follows. If m_i is even, we take a curve of genus $g_i = -m_i/2$ such that there exists a differential η_i on X_i with a unique zero of order $-m_i - 2$ at z_0 . If m_i is odd, we attach a curve of genus $g_i = (-m_i + 1)/2$ such that there exists a differential η_i on X_i with a zero of order $-m_i - 2$ at z_i and a simple zero elsewhere. Let X be the resulting stable curve.

The collection of $\eta_0, \eta_1, \dots, \eta_n$ defines a twisted differential η of type $(m_0, 1, \dots, 1)$ on X , where the number of ones is equal to the number d of odd negative m_i . Consider the level graph $\bar{\Gamma}$ on X such that $X_1 \asymp \dots \asymp X_n \succ X_0$. Given the hypothesis that η_0 has zero residues at z_1, \dots, z_n , the twisted differential η is compatible with $\bar{\Gamma}$. Hence by Theorem 1.3, all stable pointed differentials associated to such η are in the incidence variety compactification of the stratum $\mathbb{P}\Omega\mathcal{M}_g(m_0, 1, \dots, 1)$, where $g = \sum_{i=1}^n \lfloor \frac{-m_i+1}{2} \rfloor$.

Note that

$$\dim \mathbb{P}\Omega\mathcal{M}_g(m_0, 1, \dots, 1) = 2g + d - 1.$$

On the other hand, the dimension of the space of stable pointed differentials associated to the twisted differentials we constructed, with the projectivization of η_0 on X_0 fixed, has dimension

$$-1 + \sum_{i>0, m_i \text{ even}} 2g_i + \sum_{i>0, m_i \text{ odd}} (2g_i + 1) = 2g + d - 1,$$

where the -1 results from simultaneous projectivization of the differentials on the top level components. Hence it has the same dimension as the stratum, contradicting the fact that the boundary of a closed variety has smaller dimension compared to that of the interior. \square

Remark 3.7. The lemma can also be proved directly in two ways. If ω is such a differential with no residue, then $\omega = df$ for some meromorphic function f on \mathbb{P}^1 . Regarding f as a map from \mathbb{P}^1 to \mathbb{P}^1 , the unique zero of ω corresponds to the unique ramification point of f over $\mathbb{P}^1 \setminus \{\infty\}$, and the n poles of ω map to ∞ . It follows from the Riemann-Hurwitz formula that $2d - 2 = (d - 1) + (d - n)$ where d is the degree of f , which contradicts that $n \geq 2$. Alternatively if ω has no residue, then the flat geometric representation of ω in the sense of [Boi15b] has no saddle connections, hence the surface would be a wedge sum of n spheres attached at the unique zero, leading to a contradiction.

The above lemma yields a useful criterion when we apply the global residue condition to twisted differentials on a stable curve with a rational component.

Corollary 3.8. *Suppose $\mu = (m_1, \dots, m_n)$ is a holomorphic type, i.e. all m_i are positive. Let (X, z_1, \dots, z_n) be a stable curve with an irreducible component X_v of genus zero, such that all the nodes q_1, \dots, q_k contained in X_v are separating, and $k \geq 2$. If η is a twisted differential of type μ such that η_v on X_v has a unique zero and has a pole at each q_i , then η is not compatible with any level graph on X .*

Proof. Suppose, on the contrary, that there is a level graph $\bar{\Gamma}$ on X such that η is compatible with $\bar{\Gamma}$. Since η_v has a pole at each node q_i , X_v is on a lower level compared to the component on the other branch of q_i . Since q_i is separating, the global residue condition applied to the level of X_v implies that $\text{Res}_{q_i} \eta_v = 0$ for all i , contradicting Lemma 3.6. \square

Next we describe a relationship between pointed stable differentials and compatible level graphs. An irreducible component X_v of X is called a *non-strict local minimum* if for every component X_u intersecting X_v , the inequality $X_u \preceq X_v$ holds.

Lemma 3.9. *Let $(X, \omega, z_1, \dots, z_n)$ be a pointed stable differential and suppose that $g \neq 1$ or $n \neq 0$. If an irreducible component X_v of X is a (non-strict) local minimum in a level graph $\bar{\Gamma}$, then any twisted differential η compatible with $\bar{\Gamma}$ that agrees with ω on the top level components, has at least one zero or marked point at a smooth point of X_v .*

Proof. Since X_v is a (non-strict) local minimum, the differential η_v has a pole at any node lying on X_v . If η_v has no marked zero in the smooth locus of X_v , then $X_v \cong \mathbb{P}^1$, because the only Riemann surfaces on which there exist a meromorphic differential without zeros are \mathbb{P}^1 and a genus 1 curve — and in the latter case the differential must be holomorphic, with no zeroes or poles, and thus X_v must equal to X , which is the excluded case. In that case η_v either has two simple poles or a single double pole; hence for the curve to be stable, X_v must contain at least one marked point. \square

This lemma implies the following useful result.

Corollary 3.10. *For $g \geq 1$ any pointed stable differential (X, ω, z) in the incidence variety compactification $\mathbb{P}\Omega\bar{\mathcal{M}}_{g,1}^{\text{inc}}(2g-2)$ is compatible with a level graph that has a unique local minimum, which is the irreducible component of X containing the marked point z .*

We conclude this section by the following lemma for later use. Recall the map π_2 defined in (2.8) from the incidence variety compactification to the Deligne-Mumford compactification.

Lemma 3.11. *Let S be an open boundary stratum of $\bar{\mathcal{M}}_{g,n}$ parameterizing curves with a fixed dual graph. Let \tilde{S} be the boundary stratum corresponding to the dual graph obtained from the dual graph of S by adding a loop at a vertex v and decreasing the geometric genus of X_v by one (i.e., adding a non-separating node q for an irreducible component X_v of curves in S). Then the dimension of $\pi_2^{-1}(\tilde{S})$ in the incidence variety compactification of any stratum of differentials is either strictly smaller than the dimension of $\pi_2^{-1}(S)$ or both preimages are empty.*

Proof. To prove this result it suffices to show that $\pi_2^{-1}(\tilde{S})$ is contained in the closure of $\pi_2^{-1}(S)$. Let $(X, \omega, z_1, \dots, z_n)$ be a stable pointed differential in $\pi_2^{-1}(\tilde{S})$. Choose an associated twisted differential η in the sense of Definition 1.6. Since the newly added node q is an internal node, η has simple poles with opposite residues at the two branches of q by conditions (1), (2), and (3). Then one can locally smooth out q to obtain a family of twisted differentials η_t , such that the stable differential associated to

the general η_t lies in $\pi_2^{-1}(S)$. This operation of locally smoothing a simple polar node can be performed by classical plumbing (see Lemma 4.4). \square

3.3. The incidence variety compactification of $\mathbb{P}\Omega\mathcal{M}_2(2)$. In this section we work out in detail the irreducible components of the boundary $\partial\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$ in the incidence variety compactification of $\mathbb{P}\Omega\mathcal{M}_2(2)$. This stratum is connected. Moreover, π_2 maps it to the locus of Weierstraß points in $\mathcal{M}_{2,1}$. The closure of this stratum was described already in [Gen18, Section 6] by using admissible double covers. Here we match these results and demonstrate how our current machinery works. Coordinates around part of this boundary were described in [Bai07, Section 6.6].

Proposition 3.12. *The boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$ has three irreducible components, given by the closures of the loci (I), (II), and (III) defined below.*

The strategy of the proof is as follows. Recall that the boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$ is divisorial (see the beginning of Section 3.2), hence its irreducible components are two-dimensional. It thus suffices to locate all two-dimensional boundary strata of $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$. We perform the search according to the dimension of the boundary strata in $\overline{\mathcal{M}}_{2,1}$.

We first determine the π_2 -preimage in $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$ of every open divisorial stratum of $\overline{\mathcal{M}}_{2,1}$. Let X be a stable curve in the stratum Δ_0° , see Figure 7. Since X is irreducible,





FIGURE 7. A curve parameterized by Δ_0° and its level graph

there is no order to consider on the dual graph. Given a twisted differential η , the associated stable differential ω coincides with η , which is not identically zero on X . Such a differential has a double zero and two simple poles on the normalization of X , which is of genus one. Thus the π_2 -preimage of Δ_0° in $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}$ can be identified with the locus

$$(I) := \mathbb{P}\Omega\mathcal{M}_{1,\{3\}}^{\text{inc}}(2, -1, -1),$$

where the two simple poles are not ordered. This is the locus of elliptic curves E with three marked points p_1, p_2, p_3 such that $2p_1 = p_2 + p_3$. It is irreducible according to [Boi15a]. The locus (I) is two-dimensional, and hence it gives an irreducible component of the boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$.

In the open boundary divisor Δ_1° , a stable curve X is the union of two elliptic curves X_1 and X_2 intersecting at q , where $z \in X_2$. Suppose η is a twisted differential compatible with a level graph $\overline{\Gamma}$ on X . Then η_2 on X_2 has a unique zero of order two at z , hence η_2 has a double pole at q^- , and the component X_2 is of lower level than X_1 in $\overline{\Gamma}$, see Figure 8. Consequently η_1 on X_1 is holomorphic and nowhere vanishing. The global residue condition requires that $\text{Res}_{q^-} \eta_2 = 0$, which follows from the residue theorem for η_2 on X_2 . We have thus verified that in the above setting η is

FIGURE 8. A curve parameterized by Δ_1° and its level graph

compatible with $\bar{\Gamma}$. Such (X, η) is equivalent to the collection of $(X_1, \eta_1, q^+) \in \Omega\mathcal{M}_{1,1}$ and $(X_2, \eta_2, z, q^-) \in \Omega\mathcal{M}_{1,2}^{\text{inc}}(2, -2)$. Suppose ω is the associated stable differential of η . Then $\omega|_{X_1} = \eta_1$ and ω is identically zero on the lower level component X_2 . In other words, the associated pointed stable differential (X, ω, z) consists of (X_1, η_1, q^+) and (X_2, z, q^-) , which forgets η_2 compared to (X, η) . In this sense we can identify the π_2 -preimage of Δ_1° in $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}$ with the locus

$$(II) := \mathbb{P}\Omega\mathcal{M}_{1,1} \times \pi_2(\mathbb{P}\Omega\mathcal{M}_{1,2}^{\text{inc}}(2, -2)).$$

Since we are working on the open part Δ_1° of the boundary divisor, the points z and q^- are not allowed to coincide (the degeneration when z and q^- coincide corresponds to stable curves that have three irreducible components, as depicted in figure 11, and this case will be treated separately below). Thus the second factor is the locus in $\mathcal{M}_{1,2}$ where z and q^- differ by a non-zero two-torsion point. According to the classification of [Boi15a], this locus is irreducible. Together with the fact that $\mathbb{P}\Omega\mathcal{M}_{1,1}$ is irreducible, this implies that the locus (II) is irreducible and two-dimensional, and hence it gives an irreducible component of the boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$.

We now study the π_2 -preimages in $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$ of the codimension two boundary strata of $\overline{\mathcal{M}}_{2,1}$. Consider the locus of irreducible stable curves with two nodes, see Figure 9. Since this stratum is obtained by adding a loop to the dual graph of curves




FIGURE 9. An irreducible curve with two nodes

in Δ_0° , by Lemma 3.11 its π_2 -preimage has higher codimension in $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$, hence does not give a boundary component.

Next, consider the locus of stable curves as represented in Figure 10, where one component is an elliptic curve and the other component is a rational nodal curve. The marked point z is contained in one of the two components. Clearly both dual graphs are obtained by adding a loop to the dual graph of curves in Δ_1° . Hence Lemma 3.11 implies that the π_2 -preimages of these strata do not give boundary components of $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$.

Another codimension two boundary stratum of $\overline{\mathcal{M}}_{2,1}$ parameterizes stable pointed curves that have a rational bridge X_3 connecting two elliptic curves X_1 and X_2 , where



FIGURE 10. An elliptic curve union a rational nodal curve and the level graph

the marked point z lies on X_3 to ensure stability of X , see Figure 11. It follows from Corollary 3.8 that this stratum does not intersect the projection of $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$ under π_2 .




FIGURE 11. Two elliptic curves connected by a pointed rational curve

The only remaining codimension two boundary stratum of $\overline{\mathcal{M}}_{2,1}$ parameterizes curves consisting of two irreducible components X_1 and X_2 intersecting at two points, where X_1 has genus one and X_2 has genus zero, and the marked point z is contained in X_2 to ensure stability of X , see Figure 12.




FIGURE 12. Two curves intersecting at two nodes and the level graph

Suppose η is a twisted differential compatible with a level graph $\overline{\Gamma}$ on X . It follows from Corollary 3.10 that $X_1 \succ X_2$. Since η_1 on X_1 has no zeros, it is holomorphic and nowhere vanishing, and then η_2 has a double pole at each of the two nodes with a double zero at z . In this case the global residue condition requires that $\text{Res}_{q_1^-} \eta_2 + \text{Res}_{q_2^-} \eta_2 = 0$, which follows from the residue theorem. The associated stable differential ω is equal to η_1 on X_1 , and ω on X_2 is identically zero. Hence the locus of such (X, ω, z) in $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$ can be identified with

$$(III) := \mathbb{P}\Omega\mathcal{M}_{1,2} \times \mathcal{M}_{0,3},$$

where the second factor corresponds to the one-point space $\pi_2(\mathbb{P}\Omega\mathcal{M}_{0,\{3\}}^{\text{inc}}(2, -2, -2))$. Since the locus (III) is clearly irreducible and two-dimensional, it gives an irreducible component of the boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$.

It remains to consider the boundary strata of $\overline{\mathcal{M}}_{2,1}$ that have codimension greater or equal to three. If there exists a boundary component of $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$ mapping to such a stratum, then the fibers of π_2 over this stratum have positive dimension. By Lemma 2.3,

for this to happen, the graph $\bar{\Gamma}_{=0}$ must have at least two connected components. In our case, the components of top level would have to be of genus at least one. Hence such a stratum would have to parameterize curves with two irreducible components of (arithmetic) genus at least one, neither of them containing the marked point z . The only possibility is that such curves are degenerations of the curves in Figure 11. Such degenerations can happen only if (at least) one of the two elliptic components degenerates to a rational nodal curve. However, in that case Corollary 3.8 still applies, hence the locus of such curves is disjoint from the image of $\mathbb{P}\Omega\overline{\mathcal{M}}_{2,1}^{\text{inc}}(2)$ under π_2 . This completes the proof of Proposition 3.12.

3.4. The incidence variety compactification of $\mathbb{P}\Omega\mathcal{M}_3(4)$. The stratum $\mathbb{P}\Omega\mathcal{M}_3(4)$ has two irreducible components $\mathbb{P}\Omega\mathcal{M}_3(4)^{\text{hyp}}$ and $\mathbb{P}\Omega\mathcal{M}_3(4)^{\text{odd}}$ (see [KZ03]). The hyperelliptic component parameterizes hyperelliptic curves with a marked Weierstraß point. The odd-spin component parameterizes nonhyperelliptic curves whose canonical embeddings (as plane quartics) have a hyperflex. Below we will describe the closure of the entire stratum. The closure of each connected component will follow from the results in the forthcoming work [BCGGM].

For a pointed stable differential (X, ω, z) in the boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$, let $\bar{\Gamma}$ be a level graph on X and η be a compatible twisted differential such that the associated pointed stable differential is (X, ω, z) . By Corollary 3.10, $\bar{\Gamma}$ has a unique minimal vertex, and the corresponding irreducible component of X contains the marked point z .

We first determine the π_2 -preimages in $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$ of the open boundary divisors of $\overline{\mathcal{M}}_{3,1}$. The computation over Δ_0° is similar to the previous case, and we obtain the corresponding locus as

$$(i) := \mathbb{P}\Omega\mathcal{M}_{2,\{3\}}^{\text{inc}}(4, -1, -1),$$

where the two simple poles are unordered. By [Boi15a, Theorem 1.2], this locus has two connected components that can be distinguished by their parity. Both these components are four-dimensional and therefore each of them is an irreducible component of the boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$. In fact, by the discussion in [Gen18, Section 4] each component of $\mathbb{P}\Omega\mathcal{M}_{2,\{3\}}^{\text{inc}}(4, -1, -1)$ is a boundary component of the connected component of $\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$ of the same parity.

For $i = 1, 2$, denote by $\Delta_{1,i}^\circ$ the open boundary divisor of $\overline{\mathcal{M}}_{3,1}$ parameterizing nodal unions of a genus one curve with a genus two curve, where the marked point is contained in the genus i component, respectively. By Lemma 3.9, the component containing the marked point is of lower level compared to the unmarked component in $\bar{\Gamma}$. As in the previous section, we obtain that

$$(ii) := \pi_2^{-1}(\Delta_{1,1}^\circ) = \mathbb{P}\Omega\mathcal{M}_{2,1}^{\text{inc}}(2) \times \pi_2(\mathbb{P}\Omega\mathcal{M}_{1,2}^{\text{inc}}(4, -4)),$$

$$(iii) := \pi_2^{-1}(\Delta_{1,2}^\circ) = \mathbb{P}\Omega\mathcal{M}_{1,1} \times \pi_2(\mathbb{P}\Omega\mathcal{M}_{2,2}^{\text{inc}}(4, -2)),$$

both of which are again purely four-dimensional. By [Boi15a], each of these two loci has two connected components: a hyperelliptic and a non-hyperelliptic one, and each of these is then an irreducible component of the boundary of the corresponding component of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$.

Next we investigate the situation over the codimension two boundary strata of $\overline{\mathcal{M}}_{3,1}$. By Lemma 3.11, the π_2 -preimage of the locus of irreducible curves with two nodes has codimension bigger than one, hence it does not provide a boundary component of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$.

Let $\Theta_0 \subset \overline{\mathcal{M}}_{3,1}$ denote the locus of curves with two components X_0 and X_2 intersecting at two nodes, where $X_0 \cong \mathbb{P}^1$, X_2 has genus two, and X_0 contains the marked point. In this case $X_2 \succ X_0$ in $\overline{\Gamma}$, hence the stable differential ω is identically zero on X_0 . For ω on X_2 , it is given by η_2 , and there are two possibilities. If η_2 has simple zeros at both nodes, then η_0 on X_0 has triple poles at both nodes, and hence (X, ω, z) is parameterized by the irreducible locus

$$(iv) := \mathbb{P}\Omega\mathcal{M}_{2,\{2\}}^{\text{inc}}(1, 1) \times \pi_2(\mathbb{P}\Omega\mathcal{M}_{0,\{3\}}^{\text{inc}}(4, -3, -3)).$$

On the other hand, if η_2 has a double zero at one node and is regular at the other, then (X, ω, z) is parameterized by the irreducible locus

$$(v) := \mathbb{P}\Omega\mathcal{M}_{2,2}^{\text{inc}}(2, 0) \times \pi_2(\mathbb{P}\Omega\mathcal{M}_{0,3}^{\text{inc}}(4, -4, -2)).$$

Both loci are four-dimensional irreducible, and thus

$$\pi_2^{-1}(\Theta_0) = (iv) \cup (v),$$

providing (irreducible) boundary components of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$.

Let $\Theta_1 \subset \overline{\mathcal{M}}_{3,1}$ denote the locus of curves with two components X_1 and X_2 intersecting at two nodes, where both components have genus one and X_1 contains the marked point. A similar analysis as in the case of Θ_0 implies that the pointed stable differential (X, ω, z) in $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4) \cap \pi_2^{-1}(\Theta_1)$ then lies in the locus

$$(vi) := \mathbb{P}\Omega\mathcal{M}_{1,\{2\}} \times \pi_2(\mathbb{P}\Omega\mathcal{M}_{1,\{3\}}^{\text{inc}}(4, -2, -2)),$$

which has two irreducible components, both of dimension four, hence providing boundary components of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$. As remarked in [Gen18, Section 7], each of these components two components lies in the boundary of both components of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$.

For the codimension two boundary strata of $\overline{\mathcal{M}}_{3,1}$ parameterizing curves with one separating node and one non-separating node, they are contained in the closure of $\Delta_{1,i}^\circ$ for $i = 1, 2$, by pinching a non-separating loop on an irreducible component of curves in $\Delta_{1,i}^\circ$. This is the operation described by Lemma 3.11, and thus the corresponding boundary loci in $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$ are contained in (ii) and (iii) defined above. Therefore, the π_2 -preimages of such codimension two boundary strata do not give rise to any boundary components of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$.

Let us consider the other codimension two boundary strata of $\overline{\mathcal{M}}_{3,1}$ parameterizing curves with two separating nodes. There are several possibilities. If all three components X_1 , X_2 , and X_3 of the curve X have genus one and the marked point lies in the middle component X_2 , then the twisted differential η lies in the locus contained in

$$\Omega\mathcal{M}_{1,1} \times \Omega\mathcal{M}_{1,\{3\}}^{\text{inc}}(4, -2, -2) \times \Omega\mathcal{M}_{1,1},$$

which is cut out by the global residue condition that η_2 has zero residues at both nodes on the middle component X_2 . Here we encounter the situation that the map π_2 is

no longer finite. In this case the full order is $X_1 \asymp X_3 \succ X_2$. Since X_1 and X_3 are both of top level, by Lemma 2.3 the fibers of π_2 are one-dimensional, which record the projectivization of the pair of scaling factors for η_1 and η_3 . It follows that the associated pointed stable differential (X, ω, z) lies in the codimension one locus

$$(vii) \subsetneq \mathbb{P}(\Omega\mathcal{M}_{1,1} \times \Omega\mathcal{M}_{1,1}) \times \pi_2(\Omega\mathcal{M}_{1,\{3\}}^{\text{inc}}(4, -2, -2))$$

given by the condition that η_2 has zero residues. By the results of [Boi15b] it follows that the locus (vii) is non-empty, and thus its codimension in this product is equal to one. The locus (vii) is thus purely four-dimensional; the number of its irreducible components is not known, but each such irreducible component gives a boundary component of $\mathbb{P}\Omega\mathcal{M}_{3,1}^{\text{inc}}(4)$.

On the other hand if the marked point z lies in X_1 , then the full order is given by $X_3 \succ X_2 \succ X_1$. Since X_3 is the unique irreducible component of top level, the map π_2 is finite. The twisted differential η lies in the locus

$$\Omega\mathcal{M}_{1,1} \times \Omega\mathcal{M}_{1,2}^{\text{inc}}(2, -2) \times \Omega\mathcal{M}_{1,2}^{\text{inc}}(4, -4)$$

and the associated pointed stable differential lies in the locus

$$\mathbb{P}\Omega\mathcal{M}_{1,1} \times \pi_2(\Omega\mathcal{M}_{1,2}^{\text{inc}}(2, -2)) \times \pi_2(\Omega\mathcal{M}_{1,2}^{\text{inc}}(4, -4)),$$

which is three-dimensional, and thus not a boundary component of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$.

Suppose now that one of the three irreducible components of X has genus zero. By stability of X , the \mathbb{P}^1 component must be the middle component, and it contains the marked point z . It follows from Corollary 3.8 that this locus is disjoint with the image of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$ under π_2 . This completes our analysis over the codimension two boundary strata of $\overline{\mathcal{M}}_{3,1}$.

We now investigate over which boundary strata of $\overline{\mathcal{M}}_{3,1}$ of codimension greater or equal to three there may be boundary components of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$. As there are numerous cases, we first make some general observations. If a boundary stratum of $\overline{\mathcal{M}}_{3,1}$ parameterizes stable curves that have an internal node, then by Lemma 3.11, its π_2 -preimage does not give rise to a boundary component of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$. Therefore, it suffices to consider the case when all irreducible components of the stable curves are smooth. Furthermore, if the π_2 -preimage of a boundary stratum of $\overline{\mathcal{M}}_{3,1}$ of codimension greater or equal to three gives a boundary component of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$, then the fibers of π_2 over this stratum have positive dimension.

Let us first determine which codimension three strata of the boundary of $\overline{\mathcal{M}}_{3,1}$ are contained in $\pi_2(\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4))$. As before, Lemma 3.11 allows us to assume that no edge of the dual graph Γ is a loop. Moreover, if $h^1(\Gamma) \geq 3$, the stratum has codimension at least 4, and can be discarded. The case of $h^1(\Gamma) = 0$, i.e. the Γ being a tree, is also easily ruled out. If $h^1(\Gamma) = 1$, then Γ must be the triangle, with two curves of genus one and one \mathbb{P}^1 containing the marked point. In this case, the admissible twisted differentials gives a non-trivial torsion condition on the marked points on one elliptic curve. Hence the whole stratum is not in the image $\pi_2(\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4))$. Finally, if $h^1(\Gamma) = 2$, the only possibility is the locus parameterizing curves with two irreducible

components intersecting at three nodes, where one component has genus one, and the other is \mathbb{P}^1 , containing the marked point z . Moreover, the \mathbb{P}^1 must be of lower level in $\bar{\Gamma}$. The global residue condition on η then holds automatically by the residue theorem. In this case the associated pointed stable differentials are parameterized by

$$(viii) := \mathbb{P}\Omega\mathcal{M}_{1,3} \times \pi_2(\mathbb{P}\Omega\mathcal{M}_{0,\{4\}}^{\text{inc}}(4, -2, -2, -2)).$$

This locus is irreducible and four-dimensional, hence it gives an irreducible boundary component of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$.

Let us now look at the cases when π_2 has positive dimensional fibers. According to Lemma 2.3 the level graph $\bar{\Gamma}$ has at least two top level components. Both are smooth and of genus at least one. If some two top level components are connected by a node, the presence of this node decreases the codimension of such a locus by one, and thus such a locus must be properly contained in a higher-dimensional component of the boundary. Moreover by Lemma 3.9, the unique minimal component X_v of $\bar{\Gamma}$ contains the marked point z . If X_v has genus at least one, then the curve X is of compact type (which, recall, means that the dual graph is a tree). The only such stable curve X is given by a \mathbb{P}^1 component, denoted by X_3 , intersecting three elliptic components X_1 , X_2 , and X_v , with the full order given by $X_1 \asymp X_2 \succ X_3 \succ X_v$. Then a compatible twisted differential η on $X_3 \cong \mathbb{P}^1$ would have a double zero at the node joining X_v and two double poles with zero residues at the other two nodes, which is impossible by Lemma 3.6.

Now assume that the minimal component X_v has genus zero. Since X_v contains z , in order to be stable, X_v must have at least two nodes. If it has precisely two nodes, then $X \setminus X_v$ has to be connected, for otherwise the global residue condition would imply that η_v has zero residues at the two nodes and a unique zero at z , which is impossible by Lemma 3.6. In this case the only possibility of such X and $\bar{\Gamma}$ is $X_1 \asymp X_2 \succ X_v$, where X_1 and X_2 are of genus one, and $X_v \cong \mathbb{P}^1$. Then there does not exist a compatible twisted differential η , for otherwise η_1 on the elliptic component X_1 would have a simple pole at the node joining X_2 and a simple zero at the node joining X_v , impossible.

Now consider the case when $X_v \cong \mathbb{P}^1$ contains more than two nodes. If there are three top level components, all of them have to be elliptic curves and have to intersect X_v . Then the global residue condition implies that any compatible twisted differential η has zero residues at the poles of η_v on X_v . By Lemma 3.6 such η does not exist. If X has only two top level components, then both of them have to be elliptic curves, and the dual graph of X contains a circle. Hence all but two of the nodes contained in X_v are separating. In order for X to be stable, at each separating node the attached curve must be of genus one. Thus there can be only one separating node. In this case the only possibility is that X_v intersects X_1 at one point and X_v intersects X_2 at two points, where both X_1 and X_2 are of genus one. Let S be the locus in $\Omega\mathcal{M}_{0,\{4\}}^{\text{inc}}(4, -2, -2, -2)$ parameterizing meromorphic differentials whose residues are zero at the first pole. In this case the twisted differentials are parameterized by the locus

$$\Omega\mathcal{M}_{1,1} \times \Omega\mathcal{M}_{1,2} \times S.$$

The associated pointed stable differentials (X, ω, z) are parameterized by

$$(ix) := \mathbb{P}(\Omega\mathcal{M}_{1,1} \times \Omega\mathcal{M}_{1,2}) \times \pi_2(S),$$

which is four-dimensional. We claim that this locus is connected, and thus it gives an irreducible component of the boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$. Indeed, a differential in $\Omega\mathcal{M}_{0,\{4\}}^{\text{inc}}(4, -2, -2, -2)$ can be written in the form $\omega = \frac{dz}{z^2(z-1)^2(z-a)^2}$ with $a \in \mathbb{C} \setminus \{0, 1\}$. The residue of ω at 0 is $\text{Res}_0(\omega) = \frac{2(a+1)}{a^3}$, which is zero if and only if $a = -1$. This shows that S and hence (ix) is connected.

In summary, we have proved the following result.

Proposition 3.13. *The boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$ is the union of the closures of the loci (i), (ii), (iii), (iv), (v), (vi), (vii), (viii), and (ix) defined above. Furthermore, for each of these cases except (vii), the irreducible components of the corresponding locus are enumerated above, so that in total we see that the number of irreducible components of the boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{3,1}^{\text{inc}}(4)$ is equal to 12 plus the number of irreducible components of the locus (vii). Moreover, the loci (vii) and (ix) are contracted under the map π_2 .*

We remark that in [Gen18] the third author analyzed the incidence variety compactification of $\mathbb{P}\Omega\mathcal{M}_{3,1}^{\text{inc}}(4)$ over the boundary divisors Δ_0° and Δ_1° of $\overline{\mathcal{M}}_3$. In [Che17] the second author characterized the Deligne-Mumford compactification of $\mathbb{P}\Omega\mathcal{M}_{3,1}(4)$ in $\overline{\mathcal{M}}_{3,1}$ for curves with at most two nodes. In [FP16] Farkas and Pandharipande also studied the Deligne-Mumford compactification of $\mathbb{P}\Omega\mathcal{M}_{3,1}(4)$ for several cases of stable curves with a fixed dual graph. In [Hu17] Hu obtained an explicit modular form defining the locus of hyperflexes in $\overline{\mathcal{M}}_3$, that is, the image of $\mathbb{P}\Omega\mathcal{M}_3(4)^{\text{odd}}$ in $\overline{\mathcal{M}}_3$.

3.5. Some strata of meromorphic differentials in genus one. In this section we describe the boundary of the incidence variety compactification of $\Omega\mathcal{M}_1(m, -m)$ and $\Omega\mathcal{M}_1(m_1, m_2, m_3)$ in genus one.

First consider the stratum $\mathbb{P}\Omega\mathcal{M}_1(m, -m)$ for $m \geq 2$. If a stable curve in $\overline{\mathcal{M}}_{1,2}$ has a separating node, then it does not lie in the image $\pi_2(\mathbb{P}\Omega\overline{\mathcal{M}}_{1,\{2\}}^{\text{inc}}(m, -m))$. Otherwise the subcurve on one side of the separating node would have arithmetic genus zero, and by stability it would have to contain both marked points. Then a compatible twisted differential on this component would have a zero and a pole at the two marked points, respectively, and have another pole at the node, where both poles would have zero residues by the global residue condition and the residue theorem. But this is impossible by Lemma 3.6. Thus we only need to consider the open boundary divisor Δ_0° of $\overline{\mathcal{M}}_{1,2}$ parameterizing irreducible rational nodal curves and the locus Θ parameterizing two \mathbb{P}^1 components intersecting at two nodes, each containing a marked point.

The π_2 -preimage of Δ_0° in $\mathbb{P}\Omega\overline{\mathcal{M}}_{1,\{2\}}^{\text{inc}}(m, -m)$ can be identified with the locus

$$(j) \subset \mathbb{P}\Omega\mathcal{M}_{0,\{4\}}^{\text{inc}}(m, -1, -1, -m),$$

where the two simple poles are unordered, and the meromorphic differentials parameterized in (j) has zero residues at the pole of order m . Equivalently, (j) parameterizes marked rational curves $(\mathbb{P}^1, z, p, q^+, q^-)$ such that the degree m cover from \mathbb{P}^1 to \mathbb{P}^1 induced by $mz - mp$ contains q^+ and q^- in the same fiber.

Now we consider the π_2 -preimage of Θ in $\mathbb{P}\Omega\overline{\mathcal{M}}_{1,\{2\}}^{\text{inc}}(m, -m)$. Suppose a pointed stable differential (X, ω, z, p) lies in the π_2 -preimage of Θ , associated to a twisted differential η compatible with a level graph $\overline{\Gamma}$. The irreducible component of X containing

the marked zero z must be of lower level in $\bar{\Gamma}$. Since there is a marked pole on the top level component, there is no global residue condition in this case. Suppose the zeros of η on the top level component are of orders a and $m - 2 - a$ at the two nodes. Then such (X, η) is parameterized by

$$\Omega\mathcal{M}_{0,3}^{\text{inc}}(a, m - 2 - a, -m) \times \Omega\mathcal{M}_{0,3}^{\text{inc}}(m, -2 - a, a - m).$$

The associated pointed stable differential (X, ω, z, p) is parameterized by the locus

$$(\text{jj}_a) := \mathbb{P}\Omega\mathcal{M}_{0,3}^{\text{inc}}(a, m - 2 - a, -m) \times \mathcal{M}_{0,3},$$

which is a single point.

In summary, we have obtained the following result.

Proposition 3.14. *For $m \geq 2$ the boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{1,2}^{\text{inc}}(m, -m)$ consists of the union of the loci (j) and (jj_a) for $a = 0, \dots, \lfloor \frac{m-2}{2} \rfloor$ defined above.*

Next we study the strata $\Omega\mathcal{M}_1(m_1, m_2, m_3)$, where $m_1 \geq m_2 > 0$, $m_3 < 0$ and $m_1 + m_2 + m_3 = 0$ (so in particular we must have $m_3 \leq -2$). The π_2 -preimage of Δ_0° in $\mathbb{P}\Omega\overline{\mathcal{M}}_{1,\{3\}}^{\text{inc}}(m_1, m_2, m_3)$ can be identified with the locus

$$(\text{J}) \subset \mathbb{P}\Omega\mathcal{M}_{1,\{5\}}^{\text{inc}}(m_1, m_2, m_3, -1, -1)$$

defined by the condition that the residue is zero at the pole of order m_3 .

Suppose a pointed stable differential $(X, \omega, z_1, z_2, z_3)$ lies in the π_2 -preimage of the open boundary divisor in $\overline{\mathcal{M}}_{1,3}$ parameterizing curves with a separating node. If all three marked points are on the \mathbb{P}^1 component of X , any associated twisted differential η must be holomorphic and nowhere vanishing on the elliptic component of X . We thus obtain the locus of such $(X, \omega, z_1, z_2, z_3)$ as

$$(\text{JJ}) \subset \mathbb{P}\Omega\mathcal{M}_{1,1} \times \pi_2(\mathbb{P}\Omega\mathcal{M}_{0,4}^{\text{inc}}(m_1, m_2, m_3, -2))$$

where the residue at the last pole of order two is required to be zero. The locus (JJ) is one-dimensional, hence it gives rise to boundary components of $\mathbb{P}\Omega\overline{\mathcal{M}}_{1,\{3\}}^{\text{inc}}(m_1, m_2, m_3)$.

If the \mathbb{P}^1 component contains the two marked zeros z_1 and z_2 , and the elliptic component contains the marked pole z_3 , then such $(X, \omega, z_1, z_2, z_3)$ is parameterized by the locus

$$(\text{JJJ}_{m_3}) := \mathbb{P}\Omega\mathcal{M}_{1,2}^{\text{inc}}(-m_3, m_3) \times \pi_2(\mathbb{P}\Omega\mathcal{M}_{0,3}^{\text{inc}}(m_1, m_2, m_3 - 2)).$$

(with a further symmetrization $\mathbb{P}\Omega\mathcal{M}_{0,\{3\}}^{\text{inc}}(m_1, m_2, m_3 - 2)$) if $m_1 = m_2$). If the zero of order m_i lies on the elliptic component for $i = 1$ or 2 , then the \mathbb{P}^1 component must be of higher level. Since it contains a marked pole, by Theorem 1.3 there is no global residue condition imposed on an associated twisted differential η . Hence such (X, η) are parameterized by

$$\Omega\mathcal{M}_{1,2}^{\text{inc}}(m_i, -m_i) \times \Omega\mathcal{M}_{0,3}^{\text{inc}}(m_{3-i}, m_3, -2 + m_i).$$

The associated $(X, \omega, z_1, z_2, z_3)$ are parameterized by the locus.

$$(\text{JJJ}_{m_i}) := \pi_2(\mathbb{P}\Omega\mathcal{M}_{1,2}^{\text{inc}}(m_i, -m_i)) \times \mathbb{P}\Omega\mathcal{M}_{0,3}^{\text{inc}}(m_{3-i}, m_3, -2 + m_i).$$

Now suppose $(X, \omega, z_1, z_2, z_3)$ lies in the π_2 -preimage of the locus of curves with two \mathbb{P}^1 components intersecting at two nodes. If both marked zeros lie on the same

component, suppose the zeros of η at the two nodes on the other component are of orders a and $-m_3 - 2 - a$, respectively, for $0 \leq a \leq \lfloor \frac{-m_3-2}{2} \rfloor$. The associated pointed stable differentials are parameterized by the locus

$$(\text{JV}_{m_3}^a) := \mathbb{P}\Omega\mathcal{M}_{0,3}^{\text{inc}}(a, -m_3 - 2 - a, m_3) \times \pi_2(\mathbb{P}\Omega\mathcal{M}_{0,4}^{\text{inc}}(m_1, m_2, -a - 2, a + m_3))$$

(with a further symmetrization if $-a - 2 = a + m_3$). Similarly if one component of X contains only the marked zero z_i for $i = 1$ or 2 , suppose the zero orders of η at the nodes of the other component are a and $m_i - 2 - a$, respectively, for $0 \leq a \leq \lfloor \frac{m_i-2}{2} \rfloor$. Then the associated pointed stable differentials are parameterized by the locus

$$(\text{JV}_{m_i}^a) := \mathbb{P}\Omega\mathcal{M}_{0,3}^{\text{inc}}(m_i, -a - 2, a - m_i) \times \pi_2(\mathbb{P}\Omega\mathcal{M}_{0,4}^{\text{inc}}(m_{3-i}, a, m_i - a - 2, m_3)).$$

All the loci (JJJ_{m_i}) and $(\text{JV}_{m_i}^a)$ for $i = 1, 2, 3$ are one-dimensional, hence they give boundary components of $\mathbb{P}\Omega\overline{\mathcal{M}}_{1,3}^{\text{inc}}(m_1, m_2, m_3)$.

We claim that there are no further boundary components of $\mathbb{P}\Omega\overline{\mathcal{M}}_{1,3}^{\text{inc}}(m_1, m_2, m_3)$. By Lemma 3.11 we only need to consider the cases when each irreducible component of X is smooth. Over any boundary stratum of codimension three in $\overline{\mathcal{M}}_{1,3}$, in order to obtain a boundary component of $\mathbb{P}\Omega\overline{\mathcal{M}}_{1,3}^{\text{inc}}(m_1, m_2, m_3)$, the π_2 -fibers need to be positive dimensional, hence any associated twisted differential has at least two top level components in a compatible level graph. Each such top level component needs to carry a marked pole, which is impossible because there is only one marked pole. The only remaining codimension two boundary stratum of $\overline{\mathcal{M}}_{1,3}$ parameterizes stable curves consisting of three components X_0, X_1 , and X_2 , where $X_0 \cong X_2 \cong \mathbb{P}^1$, X_1 is of genus one, X_0 intersects X_1 and X_2 each at a node, X_0 contains a marked point, X_2 contains two marked points, and X_1 is unmarked. In this case $X_1 \succ X_0$ for any compatible η in a level graph. The global residue condition implies that η on X_0 has zero residues at the poles, hence it is a meromorphic differential on \mathbb{P}^1 that has a unique zero and two poles with zero residues. But this is impossible by Lemma 3.6.

In summary, we have proved the following result.

Proposition 3.15. *The boundary of $\mathbb{P}\Omega\overline{\mathcal{M}}_{1,\{3\}}^{\text{inc}}(m_1, m_2, m_3)$ with $m_1 \geq m_2 > 0$ and $m_3 < 0$ is the union of the closures of the loci (J) , (JJ) , (JJJ_{m_i}) , and $(\text{JV}_{m_i}^a)$ for $i = 1, 2, 3$ defined above.*

3.6. Weierstraß divisors on a binary curve. In this section, we apply our main result to study the limits of Weierstraß points on so-called *binary curves*. Recall that a binary curve is a curve that consists of two smooth irreducible components intersecting in a number of nodes. Suppose X is a smooth and connected curve of genus g . Recall that a point $z_0 \in X$ is a Weierstraß point if $h^0(X, gz_0) \geq 2$. Equivalently, this is to say that there exists an effective canonical divisor $m_0z_0 + m_1z_1 + \cdots + m_nz_n$ on X such that $m_0 \geq g$. We call such a divisor a *Weierstraß divisor of type μ* , where $\mu = (m_0, \dots, m_n)$ with $m_0 \geq g$. Note that Weierstraß divisors of type μ correspond to pointed differentials in the stratum $\mathbb{P}\Omega\mathcal{M}_{g,n+1}(\mu)$.

Degenerations of Weierstraß points and divisors have been studied intensively, see e.g. [EH87] for limits of Weierstraß points on nodal curves of compact type. Since the theory of limit linear series used in [EH87] does not work in full generality for curves

of non-compact type, it is more complicated to study limits of Weierstraß points on a curve of non-compact type. As a consequence of Theorem 1.3, in what follows we give an efficient description for the degeneration of Weierstraß divisors on certain binary curves, which recovers a main case treated by [EM02].

Let $(X, \omega, z_0, \dots, z_n)$ be a stable pointed differential of genus g , where X consists of two genus one curves E_1 and E_2 attached at $g - 1$ nodes q_1, \dots, q_{g-1} . We want to study when $(X, \omega, z_0, \dots, z_n)$ is contained in the incidence variety compactification of $\mathbb{P}\Omega\mathcal{M}_{g,n+1}(\mu)$. Without loss of generality, suppose E_1 contains z_0 . Suppose $\bar{\Gamma}$ is a level graph on X and η is a compatible twisted differential such that it is associated with $(X, \omega, z_0, \dots, z_n)$. Let η_i be the restriction of η on E_i for $i = 1, 2$.

If $E_1 \succcurlyeq E_2$, then $\text{ord}_{q_j}(\eta_1) \geq -1$ for all $j = 1, \dots, g - 1$. Note that

$$\text{div}(\eta_1) = \sum_{z_i \in E_1} m_i z_i + \sum_{j=1}^{g-1} \text{ord}_{q_j}(\eta_1) \cdot q_j.$$

Since $z_0 \in E_1$, it follows that

$$\text{deg}(\text{div}(\eta_1)) \geq m_0 - (g - 1) > 0 = \text{deg } K_{E_1},$$

which contradicts that η_1 is a (possibly meromorphic) differential on the genus one curve E_1 .

Consider the remaining case $E_1 \prec E_2$ in $\bar{\Gamma}$. Then $\text{ord}_{q_j}(\eta_2) \geq 0$ for all j . Since

$$\text{div}(\eta_2) = \sum_{z_i \in E_2} m_i z_i + \sum_{j=1}^{g-1} \text{ord}_{q_j}(\eta_2) \cdot q_j,$$

in order to ensure $\text{deg}(\text{div}(\eta_2)) = 0$, the only possibility is that $\text{ord}_{q_j}(\eta_2) = 0$ for all j and that E_2 does not contain any marked point z_i . Therefore, we conclude that

$$\text{div}(\eta_1) = \sum_{i=0}^n m_i z_i - 2 \sum_{j=1}^{g-1} q_j,$$

hence necessarily $\sum_{i=0}^n m_i z_i \sim 2 \sum_{j=1}^{g-1} q_j$ on E_1 , and η_2 is holomorphic and nowhere vanishing on E_2 .

Conversely if the above linear equivalence relation holds, the global residue condition imposed on η follows from the residue theorem on E_1 . In that case, take $\omega|_{E_1} = 0$ and $\omega|_{E_2} = \eta_2$. Then $(X, \omega, z_0, \dots, z_n)$ lies in the incidence variety compactification of $\mathbb{P}\Omega\mathcal{M}_{g,n+1}(\mu)$ by Theorem 1.3.

Applying the above method, it is clear that our compactification can be used to determine the degeneration of Weierstraß divisors on stable pointed curves of any given type.

4. PLUMBING AND GLUING DIFFERENTIALS

4.1. Proof of the main theorem: conditions are necessary. First we show how to obtain a collection of meromorphic differentials associated to any boundary point of the incidence variety compactification. Suppose $f : \mathcal{X} \rightarrow \Delta$ is a family of pointed curves over a sufficiently small disk Δ whose central fiber is nodal, along with (possibly

meromorphic) differentials of type $\mu = (m_1, \dots, m_n)$ on the fibers $\mathcal{X}^* \rightarrow \Delta^*$ over the punctured disk $\Delta^* := \Delta \setminus \{0\}$. We denote by X_t the fiber of f over $t \in \Delta^*$, and by $X = X_0$ the central fiber. The marked points in the family are recorded as sections $\mathcal{Z}_1, \dots, \mathcal{Z}_n : \Delta \rightarrow \mathcal{X}$ that correspond to the zeros and poles of the differentials. The family of stable differentials on \mathcal{X} can then be viewed as a section $\mathcal{W} : \Delta^* \rightarrow f_*\omega_{\mathcal{X}^*/\Delta^*}(-\mathcal{Z}_{\text{pol}})$ where $\mathcal{Z}_{\text{pol}} = \sum_{m_i < 0} m_i \mathcal{Z}_i$ is the polar part of μ .

The result below shows that for any irreducible component X_v of the central fiber, the family of differentials \mathcal{W} over Δ^* can be rescaled by a power of t so that the limit will exist and will not be identically zero on X_v .

Lemma 4.1. *Let $(f : \mathcal{X} \rightarrow \Delta, \mathcal{W}, \mathcal{Z}_1, \dots, \mathcal{Z}_n)$ be a family of pointed differentials over a disk Δ described above. Then for every irreducible component X_v of the central fiber X , there exists a unique integer ℓ_v such that for a generic section $s : \Delta^* \rightarrow \mathcal{X}$ with $\bar{s}(0) \in X_v$ there exists a nonzero limit*

$$(4.1) \quad \eta_v := \lim_{t \rightarrow 0} t^{\ell_v} \mathcal{W}(t, s(t)) \neq 0.$$

We refer to the exponent ℓ_v in the lemma as the *scaling parameter* of the component X_v and to the η_v as the *scaling limits*. If one re-parameterizes the base disk, the scaling limits on the irreducible components with the same value of the scaling parameter will be multiplied by the same nonzero scalar.

Proof. Since \mathcal{X} is normal it is regular in codimension-one, so the local ring of $\mathcal{O}_{\mathcal{X}}$ at X_v is a discrete valuation ring (DVR) (see e.g., [AM69, Theorem 9.3]). As the fiber $f^{-1}(0)$ is reduced, t is a uniformizing parameter for this DVR. It follows that any section s of a line bundle over \mathcal{X} is of the form $t^{a_v} g$, where g is another local holomorphic section of the same line bundle over \mathcal{X} which does not vanish on X_v . In these terms a_v is just the order of vanishing of s along X_v , and ℓ_v in the claim is equal to $-a_v$. \square

We now show that the collection of these scaling limits gives a twisted differential satisfying all the conditions.

Proof of Theorem 1.3: conditions are necessary. Suppose a pointed stable differential $(X, \omega, z_1, \dots, z_n)$ lies in the incidence variety compactification of $\mathbb{P}\Omega\mathcal{M}_g(m_1, \dots, m_n)$. Take a family of pointed stable differentials over a disk such that $(X, \omega, z_1, \dots, z_n)$ is the central fiber and choose an arbitrary lifting of the family of differentials over Δ^* , defined up to scale, to an actual family of differentials. Define the collection of meromorphic differentials $\{\eta_v\}$ on the irreducible components X_v of X as the scaling limits given by Lemma 4.1. Then the scaling parameters ℓ_v can be used as a level function on the vertices of the dual graph Γ of X , and thus define a level graph $\bar{\Gamma}$. Note that multiplying the section \mathcal{W} globally by a power of t results in adding a constant to all the scaling parameters ℓ_v . To ensure that the last statement of condition (i) in Theorem 1.3 holds, we perform such a global rescaling of \mathcal{W} by a power of t so that the maximum scaling parameter (corresponding to the top components) is equal to $\ell_v = 0$, and all other scaling parameters are thus negative. From now on we fix such a choice, and show that the collection $\eta := \{\eta_v\}$ gives a twisted differential compatible with $\bar{\Gamma}$.

First, note that in the incidence variety compactification the prescribed zeros and poles are marked points, which are not allowed to collide with a node of the central

fiber. Thus the vanishing orders of η at all the marked points z_i are the same as in the family, so η satisfies condition (0).

Next we prove conditions (1) and (2). Suppose q is a node joining two irreducible components X_{v_1} and X_{v_2} of the central fiber X . Take a neighborhood U of the node q and choose local coordinates x, y such that $U \cong V(xy - t^a)$ for some $a \in \mathbb{N}$. We pick a differential ξ that generates $\omega_{\mathcal{X}/\Delta}$ and that vanishes nowhere on U , for example

$$\xi = \frac{dx}{x} = -\frac{dy}{y}.$$

To establish conditions (1) and (2), we use that the orders of vanishing and residues of \mathcal{W} in each branch can be detected by the turning numbers and periods of loops in these branches. We define two families of loops $\gamma_t^1, \gamma_t^2: [0, 2\pi] \rightarrow X_t$ by

$$\begin{aligned} \gamma_t^1(\tau) &= (re^{i\tau}, t^a/re^{i\tau}, t^a) \\ \gamma_t^2(\tau) &= (t^a/re^{i\tau}, re^{i\tau}, t^a) \end{aligned}$$

for some sufficiently small $r > 0$. The orders of vanishing are the turning numbers

$$(4.2) \quad \text{ord}_q \eta_{v_j} = \text{Turn}_{\eta_j}(\gamma_0^j) - 1,$$

where the turning number $\text{Turn}_{\eta}(\gamma)$ of a curve γ with respect to a one-form η is the degree of the Gauss map

$$G(\tau) = \frac{\eta(\gamma'(\tau))}{|\eta(\gamma'(\tau))|}.$$

As the η_i are rescalings of \mathcal{W} , we have

$$\text{Turn}_{\eta_j}(\gamma_0^j) = \lim_{t \rightarrow 0} \text{Turn}_{\mathcal{W}(t)}(\gamma_t^j).$$

For t nonzero, the curve γ_t^1 is homotopic to $-\gamma_t^2$ in an annular domain of X_t in which the form \mathcal{W} is nonzero, so γ_t^1 and γ_t^2 have opposite turning numbers. It then follows from (4.2) that

$$\text{ord}_q \eta_{v_1} + \text{ord}_q \eta_{v_2} = -2,$$

which is condition (2).

When these are both simple poles, the same argument gives that the γ_t^j have opposite periods for $t \neq 0$, and by taking a limit, we see that the γ_0^j have opposite periods. The η_{v_j} then have opposite residues at q , which is condition (1).

Alternatively, conditions (1) and (2) can be explained from the viewpoint of limit linear series (see [Che17; FP16]). The equation $xy = t^a$ gives an A_{a-1} -singularity of U at the node q , which can be resolved by blowing up and inserting a chain of $a - 1$ rational curves R_1, \dots, R_{a-1} between the two branches X_{v_1} and X_{v_2} . Since the R_i are Cartier divisors after blowing up, one can use them to twist the relative dualizing line bundle such that the limit twisted differential corresponds to a section of some twisted relative dualizing line bundle. Locally twisting by a branch at a node makes the zero or pole order increase by one in one branch and decrease by one in the other branch, hence their sum remains to be -2 as in the case of the relative dualizing line bundle before twisting. Then condition (2) follows from applying this observation to

the nodes between $X_{v_1}, R_1, \dots, R_{a-1}, X_{v_2}$ and the fact that the relative dualizing line bundle restricted to R_i has degree zero. Similarly one can see condition (1) this way, as it holds for the relative dualizing line bundle without twisting.

Finally we focus on conditions (3) and (4) in Definition 1.2, which depend on the level graph. By definition of the scaling limits, we have

$$\eta_{v_1} = \lim_{t \rightarrow 0} t^{\ell_1} \mathcal{W}(t)|_{X_{v_1}}, \quad \eta_{v_2} = \lim_{t \rightarrow 0} t^{\ell_2} \mathcal{W}(t)|_{X_{v_2}}.$$

By definition of the full order, $\ell_1 \geq \ell_2$ if and only if $v_1 \succcurlyeq v_2$. If $\eta_{v_1} = x^{-1+k} dx$ for some non-negative integer k (and hence $\eta_{v_2} = y^{-1-k} dy$), then combining these conditions implies $\ell_2 - \ell_1 = k a \geq 0$. Moreover $k = 0$ if and only if $\ell_1 = \ell_2$, namely $v_1 \asymp v_2$, as claimed in condition (3).

For the global residue condition (4), as in its statement, let Y be a connected component of $X_{>L}$ that contains no marked poles. Let q_1, \dots, q_b denote the nodes where Y meets $X_{=L}$ and q_{b+1}, \dots, q_{b+c} be the nodes where Y meets the other irreducible components of X . Since Y is a connected component of $X_{>L}$ and since all components of $X_{=L}$ have already been accounted for, it follows that $q_{b+1}^-, \dots, q_{b+c}^- \in X_{<L}$.

For each $j = 1, \dots, b+c$, let $V_{q_j}(t) \in H_1(X_t, \mathbb{Z})$ be the family of vanishing cycles, i.e., $V_{q_j}(t)$ is a simple loop on X_t for $t \neq 0$ that shrinks to the point q_j^+ as $t \rightarrow 0$. We cut the surfaces X_t along all the vanishing cycles $V_{q_j}(t)$ and obtain two subsurfaces Y_t and Z_t , which we label so that Y_t converges to Y and the (possibly disconnected) subsurface Z_t converges to the complement $X \setminus Y$. Hence Y_t is a Riemann surface with boundary $\cup_{j=1}^{b+c} V_{q_j}(t)$. We provide $V_{q_j}(t)$ with the counterclockwise orientation induced by the flat structure on Y_t . Since Y_t contains no prescribed pole, by Stokes' theorem we conclude that

$$(4.3) \quad \sum_{j=1}^{b+c} \int_{V_{q_j}(t)} \mathcal{W}(t)|_{Y_t} = 0 \quad \text{for any } t \neq 0.$$

Since $\mathcal{W}(t)$ is a family of differentials on the whole surface X_t for $t \neq 0$, we can as well consider it on Z_t and multiply by t^L to obtain

$$(4.4) \quad \sum_{j=1}^{b+c} \int_{V_{q_j}(t)} t^L \mathcal{W}(t)|_{Z_t} = 0 \quad \text{for any } t \neq 0.$$

Now we observe that $t^L \mathcal{W}(t)|_{Z_t}$ converges, as $t \rightarrow 0$, to η_v for every component $X_v \subset X_{=L}$, and converges to zero on $X_{<L}$. Since the limit of the loops $V_{q_j}(t)$ does not lie on any component of level $> L$ by construction, this means taking the limit as $t \rightarrow 0$ of (4.4) gives precisely the global residue condition (4). \square

4.2. Deformations of standard coordinates. In this section and next, we will establish a number of technical results that are needed for proving the sufficiency of the conditions in the main theorem.

The two obvious local conformal invariants of a holomorphic one-form ω defined in a small disk containing 0 are its order of vanishing $k = \text{ord}_0 \omega$ and its residue $r = \text{Res}_0 \omega$. It is somewhat less obvious that these are the only local invariants of ω . More precisely, there is a nearly canonical change of coordinates which puts ω in a standard normal

form depending only on k and r . If ω is a holomorphic one-form on a small disk, and has a zero of order k at the origin, one can take a coordinate z such that

$$z(p)^{k+1} = (k+1) \int_0^p \omega,$$

which makes $\omega = z^k dz$. For meromorphic one-forms, the residue needs to be accounted for, and the general statement is the following.

Proposition 4.2. *Given any meromorphic one-form ω on a sufficiently small disk V around 0, denote $k := \text{ord}_0 \omega$ and $r := \text{Res}_0 \omega$. Then there exists a conformal map $\phi: (\Delta_R, 0) \rightarrow (V, 0)$ defined on a disk of sufficiently small radius R such that*

$$(4.5) \quad \phi^* \omega = \begin{cases} z^k dz & \text{if } k \geq 0, \\ \frac{r}{z} dz & \text{if } k = -1, \\ \left(z^k + \frac{r}{z}\right) dz & \text{if } k < -1. \end{cases}$$

The germ of ϕ is unique up to multiplication by a $(k+1)$ st root of unity when $k > -1$, and up to multiplication by a nonzero constant if $k = -1$.

Moreover for a neighborhood U of the origin $\mathbf{0} \in \mathbb{C}^n$, let $\omega_{\mathbf{u}}$ be a family of meromorphic differentials on V that vary holomorphically with the base parameter $\mathbf{u} \in U$ and that satisfy $\text{ord}_0 \omega_{\mathbf{u}} \equiv k$. Then in some neighborhood of any point in U , one can find for some $R > 0$ a family of conformal maps $\phi_{\mathbf{u}}: (\Delta_R, 0) \rightarrow (V, 0)$, varying holomorphically with \mathbf{u} , such that $\phi_{\mathbf{u}}^* \omega_{\mathbf{u}}$ is in the standard form of (4.5).

Proof. The desired coordinates are constructed in [Str84] as a solution to a suitable ordinary differential equation. The last statement follows immediately from holomorphic dependence on parameters. \square

We call the above coordinate z a *standard coordinate* for ω .

For families of differentials such that $\text{ord}_0 \omega_{\mathbf{u}}$ is not constant, the situation is more complicated. Given a form $\omega = z^k dz$ in standard coordinates, for $k \geq 0$, a domain $U \subset \mathbb{C}^n$, and a holomorphic map $\alpha: \Delta_R \times U \rightarrow \mathbb{C}$, written as $\alpha_{\mathbf{u}}(z)$ and such that $\alpha_{\mathbf{0}}$ is identically zero, consider the family of one-forms

$$(4.6) \quad \omega_{\mathbf{u}} := \left(z^k + \frac{\alpha_{\mathbf{u}}(z)}{z} \right) dz,$$

which we call an α -deformation of ω . In this deformation, the zero of order k breaks up into a simple pole at 0 and $k+1$ nearby zeros (or k zeros if $\alpha_{\mathbf{u}}(0) \equiv 0$).

While the above standard coordinates can no longer be defined on a neighborhood of 0, we now establish that these coordinates persist on an annulus $\{R_1 < |z| < R_2\}$, where all zeros of $\omega_{\mathbf{u}}$ are contained in $\{|z| < R_1\}$.

Theorem 4.3. *Consider the annulus $A := \{z : R_1 < |z| < R_2\} \subset \Delta_R$ for some fixed $R_1 < R_2 < R$. Let $\omega := z^k dz$ be the holomorphic differential in the standard form on Δ_R for some $k \geq 0$, $\omega_{\mathbf{u}}$ an α -deformation of ω , and let $\theta_j := e^{2\pi i j / (k+1)}$ be a $(k+1)$ st root of unity. Choose a basepoint $p \in A$ and a holomorphic map $\sigma: U \rightarrow \Delta_R$ such that $\sigma(\mathbf{0}) = \theta_j p$.*

Then there exists a neighborhood $U_{\mathbf{0}} \subset U$ of $\mathbf{0}$ and a holomorphic map $\phi : A \times U_{\mathbf{0}} \rightarrow \Delta_R$, such that $\phi_{\mathbf{u}}^*(\omega_{\mathbf{u}}) = \eta_{\alpha_{\mathbf{u}}(0)}$ where

$$\eta_r = \left(z^k + \frac{r}{z} \right) dz,$$

with $\phi_{\mathbf{0}}(z) = \theta_j z$ for all $z \in A$ and with $\phi_{\mathbf{u}}(p) = \sigma(\mathbf{u})$ for all $\mathbf{u} \in U_{\mathbf{0}}$.

We remark that using $\theta_j = 1$ would be enough for our purpose here, but the freedom of choosing a different $(k+1)$ st root θ_j will be useful in our forthcoming paper [BCGGM]. Moreover, the theorem allows $\alpha_{\mathbf{u}}(0) = 0$ for all \mathbf{u} , namely, it allows us to work with deformations that do not have residues. This will be important for the procedure of merging nearby zeroes to a higher order zero, which we will develop later.

Proof. Given a holomorphic function $h = h_{\mathbf{u}}(z)$ on A depending holomorphically on the parameter $\mathbf{u} \in U$, let ϕ_h denote the holomorphic function

$$\phi_h(z) := \theta_j \cdot z \cdot e^{h(z)}.$$

We interpret the condition $\phi_h^*(\omega_{\mathbf{u}}) = \eta_{\alpha_{\mathbf{u}}(0)}$ as a differential equation involving h and α . The Implicit Function Theorem for Banach spaces (see [Lan93, Theorem XIV.2.1] or [Whi65]) will then allow us to perturb the solution $\phi_{\mathbf{0}}(z) = \theta_j z$ (with $h_{\mathbf{0}}(z) \equiv 0$) to obtain a holomorphically varying family of solutions.

Let $\mathcal{O}(A)_m$ denote the Banach space of holomorphic functions on A whose first m derivatives are uniformly bounded, equipped with the norm

$$\|f\|_m := \sum_{j=0}^m \sup_{z \in A} |f^{(j)}(z)|.$$

We calculate $\phi_h^*(\omega_{\mathbf{u}}) - \eta_{\alpha_{\mathbf{u}}(0)} = F(h, \mathbf{u})dz$, where

$$F(h, \mathbf{u}) = \left(h' + \frac{1}{z} \right) \left(z^{k+1} e^{(k+1)h} + \alpha_{\mathbf{u}}(\theta_j z e^h) \right) - z^k - \frac{\alpha_{\mathbf{u}}(0)}{z}.$$

To make sure that $F(h, \mathbf{u})$ is defined, we choose a neighborhood of the identically zero function $W \subset \mathcal{O}(A)_1$ such that for any h in W , the function ϕ_h is univalent on A , takes values in Δ_R , and 0 lies in the bounded component of the complement of $\phi_h(A)$. Note that ϕ_h is univalent on A as long as $\psi_h(z) = \phi_h(z) - z$ satisfies $|\psi_h'(z)| < 1/2$ on A , since $\phi_h(z_1) = \phi_h(z_2)$ implies $|z_1 - z_2| = |\psi_h(z_1) - \psi_h(z_2)| \leq \frac{1}{2}|z_1 - z_2|$, so $z_1 = z_2$.

We then regard F as a holomorphic map $F : W \times U \rightarrow H$, where $H \subset \mathcal{O}(A)_0$ is the subspace of functions with zero coefficient of z^{-1} in the Laurent expansion. Note that H is a closed subspace, and thus itself a Banach space, as the zero coefficient condition is equivalent to the residue being zero, and computing the integral over a path is a continuous linear map. That the image of F lies in H follows immediately from the definition of F , since for any curve γ generating the fundamental group of A , the image $\phi_h(\gamma)$ winds once around zero, so

$$\int_{\gamma} \phi_h^* \omega_{\mathbf{u}} = \int_{\phi_h(\gamma)} \omega_{\mathbf{u}} = 2\pi i \alpha_{\mathbf{u}}(0) = \int_{\gamma} \eta_{\alpha_{\mathbf{u}}(0)}.$$

It follows that F has no z^{-1} term.

To handle the initial condition, we define $G: W \times U \rightarrow H \times \mathbb{C}$ by

$$G(h, \mathbf{u}) := (F(h, \mathbf{u}), \theta_j p e^{h(p)} - \sigma(\mathbf{u})).$$

We then have $G(0, \mathbf{0}) = (0, 0)$. We wish to apply the Implicit Function Theorem to produce a neighborhood $U_{\mathbf{0}} \subset U$ of $\mathbf{0}$ and a holomorphic function $\Phi: U_{\mathbf{0}} \rightarrow W$ such that $G(\Phi(\mathbf{u}), \mathbf{u}) \equiv 0$. The desired family of holomorphic maps is then given by $\phi_{\Phi(\mathbf{u})}$. To apply the Implicit Function Theorem, it suffices to show that $D_h G_{(0, \mathbf{0})}: \mathcal{O}(A)_1 \rightarrow H \times \mathbb{C}$ (the derivative of G at $(0, \mathbf{0})$ with respect to h) is a linear Banach space homeomorphism. We first compute $D_h F_{(0, \mathbf{0})}: \mathcal{O}(A)_1 \rightarrow H$ as follows:

$$\begin{aligned} D_h F_{(0, \mathbf{0})}(h) &= \frac{d}{dt} F(th, \mathbf{0})|_{t=0} \\ &= \frac{d}{dt} \left[\left(th' + \frac{1}{z} \right) z^{k+1} e^{(k+1)th} \right]_{t=0} \\ &= z^{k+1} h' + (k+1) z^k h \\ &= (z^{k+1} h)'. \end{aligned}$$

The derivative $D_h F_{(0, \mathbf{0})}$ is thus a bounded linear operator with kernel spanned by z^{-k-1} . It also follows that

$$D_h G_{(0, \mathbf{0})}(h) = ((z^{k+1} h)', \theta_j p h(p)).$$

Define an operator $T: H \times \mathbb{C} \rightarrow \mathcal{O}(A)_1$ by

$$T(h, c) := z^{-k-1} \tilde{h},$$

where \tilde{h} is the antiderivative of h such that the initial condition $\tilde{h}(p) = \theta_j^{-1} p^k c$ holds. One checks that T and $D_h G_{(0, \mathbf{0})}$ are inverse operators, and hence $D_h G_{(0, \mathbf{0})}$ is a linear Banach space homeomorphism as desired. \square

4.3. Plumbing construction. Plumbing gives a way to deform a nodal Riemann surface to a smooth one, by plumbing the two components of the punctured neighborhood of a node (see e.g. [Wol13] for a modern treatment). Starting with a twisted differential, we would like to construct suitable meromorphic differentials on the plumbed Riemann surface.

Let us first outline the key steps involved in the plumbing construction. We want to open up a node with a holomorphic differential on the higher level component and a meromorphic differential with a pole at the node on the lower level component. When this pole has no residue, from the flat geometric point of view the plumbing is a local cut-and-paste construction (see [Gen18] and [Che17]).

When the polar node on the lower level component has a nonzero residue, it is not possible to plumb by a purely local construction, because such nonzero residues correspond to periods of curves far from the nodes under the flat metric. To deal with this issue, we first add to the differential on the higher level component a small meromorphic differential with simple poles at the nodes such that their residues are opposite to that of the poles on the lower level component. This allows us to plumb the nodes by gluing along annuli via the conformal maps constructed in Theorem 4.3. This is done in Theorem 4.5 below.

Now, when adding this meromorphic differential, a zero of the twisted differential on the higher level component may break into a number of nearby zeroes of lower order with the same total multiplicity. In Lemma 4.7 below, we describe a local operation merging these zeros back together by cutting out a neighborhood containing all these zeros and gluing in along an annulus the form $z^k dz$ supported on a small disk, again using the conformal maps constructed in Theorem 4.3. This ensures that the resulting flat surfaces are contained in the prescribed stratum.

We now set up the plumbing construction in more detail. Suppose \mathcal{X} is a family of nodal Riemann surfaces over the disc Δ with a persistent node along the section $n(t)$ together with a relative meromorphic one-form \mathcal{W} , that is, a holomorphic family of meromorphic one-forms modulo differentials pulled back from Δ . We denote by ω_t the restriction of \mathcal{W} to the fiber X_t . We want to modify $(\mathcal{X}, \mathcal{W})$ to obtain a family of smooth Riemann surfaces equipped with one-forms degenerating to the singular fiber (X_0, ω_0) . The construction is *local* near the section of nodes $n(t)$ and we will often tacitly shrink the neighborhood of $n(t)$ that we work in.

To smooth such a family, we use the standard *plumbing fixture*. Let a be a positive integer. Consider the family $\pi_a: \mathbb{V}_a \rightarrow \Delta_1$ of cylinders degenerating to a node

$$\mathbb{V}_a = \{(u, v, t) \in \Delta_1^3: uv = t^a\}$$

where the projection is given by $\pi_a(u, v, t) = t$. Let Ω be a relative meromorphic differential on \mathbb{V}_a , which we think of as a model for the plumbed family defined later. Denote by Ω_t the restriction of Ω to the fiber over t .

We first introduce some notation for the precise *plumbing setup*:

- Let $\pi^+: \mathcal{X}^+ \rightarrow \Delta_\epsilon$ and $\pi^-: \mathcal{X}^- \rightarrow \Delta_\epsilon$ be two holomorphic families of (possibly nodal) Riemann surfaces over a disk, equipped with two relative meromorphic one-forms \mathcal{W}^+ and \mathcal{W}^- . We remark that the two families are allowed to be the same for classical plumbing at a non-separating simple polar node (see Proposition 4.4 below). We denote by ω_t^+ and ω_t^- the restriction of these forms to the fibers over t .
- Let $n^\pm: \Delta_\epsilon \rightarrow \mathcal{X}^\pm$ be two holomorphic sections of these families away from the nodes of \mathcal{X}^\pm , and let $\pi: \mathcal{X} \rightarrow \Delta_\epsilon$ be the family of nodal Riemann surfaces obtained by identifying \mathcal{X}^+ and \mathcal{X}^- along n^+ and n^- to form the section of nodes n (and possibly along other sections n_t^\pm to form additional nodes that we will ignore as the plumbing construction is local at n). For $t \in \Delta_\epsilon$, we denote by

$$X_t = X_t^+ \cup X_t^- = X_t^+ \sqcup X_t^- / (n^+(t) \sim n^-(t))$$

the fiber of \mathcal{X} over t .

- For some $0 < \delta \ll 1$, let $U_\delta = \Delta_\delta^2 \times \{0\} \cap \mathbb{V}_a$ be a neighborhood of the node in the central fiber of \mathbb{V}_a , and let $D \subset X_0$ be a neighborhood of the node in the central fiber of \mathcal{X} .
- We write $\mathbb{V}_{a, \delta, \epsilon}$ for the restricted family $\mathbb{V}_a \cap (\Delta_\delta^2 \times \Delta_\epsilon) \rightarrow \Delta_\epsilon$. For some fixed $0 < \delta' < \delta$ and $0 < \epsilon < (\delta')^{2/a}$, we let $\mathcal{A} := \mathbb{V}_{a, \delta, \epsilon} \setminus \overline{\mathbb{V}_{a, \delta', \epsilon}}$ (where the bar means simply the closure). The bound on ϵ implies that \mathcal{A} is the disjoint union of two families of annuli $\mathcal{A} = \mathcal{A}^+ \cup \mathcal{A}^-$.

- Let $\mathcal{E}^\pm \subsetneq \mathcal{D}^\pm \subset \mathcal{X}^\pm$ be two families of conformal disks centered around n^\pm , so that $\mathcal{B}^\pm := \mathcal{D}^\pm \setminus \overline{\mathcal{E}^\pm}$ are two families of annuli, disjoint from the nodes. We require that the intersection of \mathcal{D}^\pm with X_0 is contained in D .

We will often omit superscripts when we take the union of two objects in the above setup, e.g. $\mathcal{E} = \mathcal{E}^+ \cup \mathcal{E}^-$.

Given the plumbing setup, we say that the family $(\mathcal{X}, \mathcal{W})$ is *plumbable with fixture* (\mathbb{V}_a, Ω) , if the following conditions hold:

- There exists a conformal isomorphism $\phi: U_\delta \rightarrow D$ such that $\phi^*\omega_0 = \Omega$.
- There exist two families of conformal isomorphisms $\Phi^\pm: \mathcal{A}^\pm \rightarrow \mathcal{B}^\pm$ such that $(\Phi^\pm)^*\mathcal{W} = \Omega$ and such that the restriction of Φ^\pm to the central fiber agrees with ϕ .

If $(\mathcal{X}, \mathcal{W})$ is plumbable with fixture (\mathbb{V}_a, Ω) , let $(\mathcal{X}', \mathcal{W}')$ be the family obtained by identifying $\mathcal{X} \setminus \overline{\mathcal{E}}$ and $\mathbb{V}_{a,\delta,\epsilon}$ along \mathcal{A}^\pm and \mathcal{B}^\pm via the maps Φ^\pm . Namely,

$$\mathcal{X}' := ((\mathcal{X} \setminus \overline{\mathcal{E}}) \sqcup \mathbb{V}_{a,\delta,\epsilon}) / (\mathcal{A} \overset{\Phi}{\sim} \mathcal{B}),$$

and \mathcal{W}' is the relative meromorphic differential induced by \mathcal{W} and Ω . In this case we say that $(\mathcal{X}', \mathcal{W}')$ is the *plumbed family with fixture* (\mathbb{V}_a, Ω) . Note that $(\mathcal{X}', \mathcal{W}')$ is smooth except for the central fiber, which is isomorphic to the nodal surface (X_0, ω_0) .

Our plumbing construction differs from other constructions in that our gluing maps are defined on a family of annuli of fixed moduli (with inner radius δ' and outer radius δ), instead of a family of growing annuli whose moduli tend to infinity. This is necessary, because Theorem 4.3, which will be used in several plumbing contexts, only constructs conformal maps on a fixed annulus.

The case of two differentials with simple poles and opposite residues is known in the literature as classical plumbing. We recall this construction here using the above notation.

Proposition 4.4 (Classical plumbing). *Suppose in the plumbing setup each of the two differentials ω_t^\pm has a simple pole at $n^\pm(t)$, with residue $r^\pm(t)$. If $r^+(t) + r^-(t) = 0$ for every t , then the family is plumbable with fixture (\mathbb{V}_1, Ω) , where*

$$\Omega_t := r^+(t) \frac{du}{u} = r^-(t) \frac{dv}{v}.$$

Proof. By working locally in the neighborhood of the pole, we transfer the problem to the setup of Proposition 4.2. We choose two holomorphically varying coordinates given by Proposition 4.2 $\psi^\pm: \Delta_\delta \rightarrow \mathcal{X}^\pm$ such that $\psi^\pm(0) = n^\pm(0)$ and such that the pullback is in the standard form:

$$(\psi_t^+)^*\omega_t^+ = r^+(t) \frac{dw}{w} \quad \text{and} \quad (\psi_t^-)^*\omega_t^- = r^-(t) \frac{dw}{w}.$$

We define the gluing map ϕ on the central fiber of $\mathbb{V}_{1,\delta,\epsilon}$ by $\phi(u, 0, 0) = \psi_0^+(u)$ and $\phi(0, v, 0) = \psi_0^-(v)$. Choose any δ' with $0 < \delta' < \delta$ and define $\mathcal{A} = \mathbb{V}_{1,\delta,\epsilon} \setminus \overline{\mathbb{V}_{1,\delta',\epsilon}}$ as in the plumbing setup. On the upper component \mathcal{A}^+ of \mathcal{A} , define the gluing map $\Phi^+(u, v, t) = \psi_t^+(u)$, and on the lower component \mathcal{A}^- , define the gluing map $\Phi^-(u, v, t) = \psi_t^-(v)$. Then these gluing maps produce the desired plumbing family. \square

We now turn to the crucial case of plumbing a zero of order $k \geq 0$ on \mathcal{X}^+ to a pole of order $k + 2$ on \mathcal{X}^- . As the pole may have a nonzero residue, we need a family of meromorphic differentials $\xi = \{\xi_t\}$ on \mathcal{X}^+ , such that ξ_t has a simple pole at the node of X_t^+ with residue opposite to that of ω_t^- . Scale ξ_t by a suitable power of t , add it to ω_t^+ , and scale ω_t^- by the same power of t . We will then plumb the modified families of differentials on \mathcal{X}^+ and \mathcal{X}^- by using the standard coordinates for the deformation, constructed in Theorem 4.3. We call ξ the *modification differential* for this plumbing procedure.

Theorem 4.5 (Higher order plumbing). *In the plumbing setup, suppose that for any t the form ω_t^+ has a zero of order $k \geq 0$ at $n^+(t)$, while ω_t^- has a pole of order $k + 2$ at $n^-(t)$ with residue $r(t)$. Let $\{\xi_t\}$ be a family of meromorphic differentials on \mathcal{X}^+ which has simple poles at $n^+(t)$ with residues $-r(t)$ and is otherwise holomorphic in a neighborhood of $n^+(t)$. Consider the family of differentials $(\mathcal{X}, \mathcal{N})$ given by*

$$(4.7) \quad \eta_t^+ = t^c(\omega_t^+ + t^b\xi_t) \text{ on } \mathcal{X}^+ \quad \text{and} \quad \eta_t^- = t^{b+c}\omega_t^- \text{ on } \mathcal{X}^-,$$

where c is a nonnegative integer and $b := a(k + 1)$ for some positive integer a , and let Ω be the family

$$\Omega_t := t^c \left(u^k - t^b \frac{r(t)}{u} \right) du = t^{b+c} \left(-v^{-k-2} + \frac{r(t)}{v} \right) dv$$

of meromorphic differentials on \mathbb{V}_a .

Then the family $(\mathcal{X}, \mathcal{N})$ is plumbable with fixture (\mathbb{V}_a, Ω) . Moreover, the scaling limit in the sense of Lemma 4.1 of the plumbed family of differentials $(\mathcal{X}', \mathcal{N}')$ on X_0^\pm is equal to ω_0^\pm .

We remark that it suffices to prove the case $c = 0$, and then the general case follows from multiplying by t^c at relevant places. The reason we include the exponent c is for later use when we apply the theorem in Section 4.4, where c and $b + c$ will be related to the scaling parameters of the two branches at the plumbed node.

Proof. As said above, we only need to consider the case $c = 0$. We want to construct the gluing maps ϕ and Φ in the definition of plumbing with prescribed fixture. By Proposition 4.2 (and multiplying the forms by -1 in the second case), we can choose two coordinates $\psi_t^\pm: \Delta_\delta \rightarrow X_t^\pm$, varying holomorphically with t in Δ_ϵ , such that $\psi_t^\pm(0) = n^\pm(t)$,

$$(\psi_t^+)^*\omega_t^+ = w^k dw, \quad \text{and} \quad (\psi_t^-)^*\omega_t^- = \left(-w^{-k-2} + \frac{r(t)}{w} \right) dw.$$

We define the gluing map ϕ on the central fiber of $\mathbb{V}_{a,\delta,\epsilon}$ by $\phi(u, 0, 0) = \psi_0^+(u)$ and $\phi(0, v, 0) = \psi_0^-(v)$. In these coordinates on the upper component, we have

$$(\psi_t^+)^*(\omega_t^+ + t^b\xi_t) = \left(w^k + \frac{\alpha_t(w)}{w} \right) dw,$$

where $\alpha_t(w)$ is holomorphic in w and t , and $\alpha_t(0) = -t^b r(t)$. Let $A = \{u : \delta_1 < |u| < \delta_2\}$ for some $0 < \delta_1 < \delta_2 < \delta$. Theorem 4.3 then gives (after possibly shrinking the

base) a family of conformal maps $\gamma_t: A \rightarrow \Delta_\delta$ such that

$$\gamma_t^* \left(w^k + \frac{\alpha_t(w)}{w} \right) dw = \left(u^k - t^b \frac{r(t)}{u} \right) du$$

The desired gluing map on the upper component \mathcal{A}^+ of $\mathcal{A} = \mathbb{V}_{a,\delta_2,\epsilon} \setminus \mathbb{V}_{a,\delta_1,\epsilon}$ is then defined by $\Phi^+(u, v, t) = \psi_t^+ \circ \gamma_t(u)$. The gluing map on the lower component \mathcal{A}^- is simply $\Phi^-(u, v, t) = \psi_t^-(v)$. By construction Φ identifies Ω and \mathcal{N} as desired. \square

To make this theorem useful in smoothing, we need a criterion describing when the modification differential ξ can be constructed globally on a Riemann surface.

Lemma 4.6. *Let $f: \mathcal{X} \rightarrow \Delta$ be a family of stable curves with sections q_j for $j = 1, \dots, n$. Suppose that $r_j(t)$ for $j = 1, \dots, n$ is a collection of holomorphic functions on Δ with $\sum_{j=1}^n r_j(t) = 0$. Then there exists a family of meromorphic differentials ξ_t on \mathcal{X} , varying holomorphically with t , with at worst simple poles at the q_j and at the nodes, satisfying the opposite-residue condition at the nodes, holomorphic outside these sections and prescribed nodes, and such that $\text{Res}_{q_j(t)}(\xi_t) = r_j(t)$ for all j and all $t \in \Delta$.*

Proof. For a single smooth Riemann surface X , the claim follows from the classical Mittag-Leffler problem (see e.g. [For91, Theorem 18.11]).

To justify the holomorphic dependence we want to find a section of $f_*\omega_{\mathcal{X}/\Delta}(\sum q_j)$ that has a prescribed image under the fiberwise linear map Res to the trivial bundle $\mathbb{C}^n \rightarrow \Delta$. Note that the set of sections with the prescribed image under Res is non-empty by the preceding paragraph, and is an affine bundle modelled on a trivial vector bundle over a disc. Such a bundle over a disc admits a holomorphic section as desired.

The proof applies to the case of stable curves as well, by treating $\omega_{\mathcal{X}/\Delta}$ as the relative dualizing sheaf. Alternatively, the stable version can be deduced from the smooth version by degenerating to the desired nodes via classical plumbing. \square

Now, when adding the modification differential ξ_t to ω_t^+ as in (4.7), a zero of ω_t^+ may in general break into many nearby zeros of lower order. In order to construct a family of differentials that remains in the initial stratum, we describe one more plumbing construction which merges these nearby zeros back into a single zero.

Our setup here is similar to the plumbing setup described above, except we will only modify a single family of Riemann surfaces. It is explicitly described as follows:

- Let $\mathcal{X} \rightarrow \Delta_\epsilon$ be a family of (possibly nodal) Riemann surfaces, and let \mathcal{W} be a family of holomorphic one-forms on \mathcal{X} which vanishes to order $k > 1$ at a smooth point p in the central fiber. We denote the restriction of \mathcal{W} to the fiber X_t by ω_t .
- Let $\phi_0: \Delta_\delta \rightarrow X_0$ be a conformal map onto a neighborhood D of p in X_0 such that $\phi_0^*\omega_0 = w^k dw$ in the standard coordinates.
- Let $\mathcal{E} \subset \mathcal{D} \subset \mathcal{X}$ be two families of conformal disks such that $\mathcal{D} \cap X_0 \subset D$, and let $\mathcal{B} = \mathcal{D} \setminus \overline{\mathcal{E}}$, a family of conformal annuli. Let \mathcal{A} be the constant family of annuli $(\Delta_\delta \setminus \overline{\Delta_{\delta'}}) \times \Delta_\epsilon \rightarrow \Delta_\epsilon$.

Given this setup, if there exists a family of conformal maps $\Phi: \mathcal{A} \rightarrow \mathcal{B}$ whose restriction to the central fiber agrees with ϕ_0 , and such that $\Phi^*\mathcal{W} = w^k dw$ for all t , then

we can construct a new family $(\mathcal{X}', \mathcal{W}')$ by gluing $\mathcal{X} \setminus \overline{\mathcal{E}}$ to $\Delta_\delta \times \Delta_\epsilon$ along the annuli via the map Φ . The central fiber (X'_0, ω'_0) is isomorphic to (X_0, ω_0) , and each ω'_t has a zero of order k . In this case we say that the resulting family $(\mathcal{X}', \mathcal{W}')$ is a *merging* of $(\mathcal{X}, \mathcal{W})$ at p .

Lemma 4.7. *Given a family $(\mathcal{X}, \mathcal{W})$ of (possibly nodal) Riemann surfaces with a family of one-forms vanishing at a smooth point $p \in X_0$ to order $k > 1$, there exists a merging of $(\mathcal{X}, \mathcal{W})$ at p .*

Proof. We provide two proofs here. The first is an application of Theorem 4.3. Extend ϕ_0 to a holomorphic family of conformal maps $\phi_t: \Delta_\delta \rightarrow X_t$. Pulling back ω_t yields

$$\phi_t^* \omega_t = (w^k + \beta_t(w))dw,$$

where β is holomorphic in t and w with $\beta_0 \equiv 0$. This is an α -deformation of $w^k dw$, where $\alpha_t(w) = w\beta_t(w)$, so $\alpha_t(0) = 0$ for all t . Fix constants $0 < \delta_1 < \delta_2 < \delta$ and let $A = \{w: \delta_1 < |w| < \delta_2\}$. Theorem 4.3 then yields a family of conformal maps $\psi_t: A \rightarrow \Delta_\delta$ such that $\psi_t^* \phi_t^* \omega_t = w^k dw$. Defining $\Phi(w, t) = \phi_t \circ \psi_t$ then provides the desired gluing map, which completes the construction.

Alternatively, we can prove the lemma without using Theorem 4.3, because the differentials involved here have no residues. Let \mathcal{W} and $\widetilde{\mathcal{W}}$ be respectively the relative differentials $(u^k + a_2(t)u^{k-2} + \dots + a_k(t)) du$ (see e.g. [KZ03, Proposition 3]) and $u^k du$ on $\mathbb{A}^1 \times \Delta$. Let $x_0(t) = (0, t)$ be the common center of masses of the zeros of these differentials in X_t . Define the maps

$$f(u, t) = \int_{x_0(t)}^{x(t)} \mathcal{W}|_{\mathbb{A}_t^1} \quad \text{and} \quad \widetilde{f}(u, t) = \int_{x_0(t)}^{x(t)} \widetilde{\mathcal{W}}|_{\mathbb{A}_t^1}.$$

Let $0 < r < R$ such that $|f(u, t)| < r$ for all zeros u of \mathcal{W} . By construction, for each t the annuli

$$A := \{r < |f(u, t)| < R\} \quad \text{and} \quad \widetilde{A} := \{r < |\widetilde{f}(u, t)| < R\}$$

are $(k+1)$ -covers (as flat surfaces defined by the corresponding differentials) of the round annulus $A_0(r, R) = \{z \in \mathbb{C} : r < |z| < R\}$, equipped with dz . Consequently, there is a biholomorphism $\Phi: \widetilde{A} \rightarrow A$ with $\Phi^* \mathcal{W} = \widetilde{\mathcal{W}}$, which provides the desired local modification. \square

4.4. Proof of the main theorem: conditions are sufficient. For future use in [BCGGM] we prove in this section a statement that is slightly stronger than the sufficiency in Theorem 1.3, since we construct a degenerating family that has not only a given twisted differential η as its scaling limit, but moreover the scaling parameters define a prescribed level graph (rather than just some level graph with which η is compatible).

Proposition 4.8. *Let $(X, \omega, z_1, \dots, z_n)$ be a pointed stable differential and let $\eta = \{\eta_v\}$ be a twisted differential on X compatible with a level graph $\overline{\Gamma}$ on X , satisfying the conditions (i), (ii), and (iii) of Theorem 1.3.*

Then there exists a family $(f: \mathcal{X} \rightarrow \Delta, \mathcal{W}, \mathcal{Z}_1, \dots, \mathcal{Z}_n)$ of pointed stable differentials over a disk Δ with parameter t , smooth outside $t = 0$, such that the set of differentials

obtained from f as scaling limits as in (4.1) coincides with η and such that its scaling parameters define precisely the level graph $\overline{\Gamma}$.

The sufficiency part of the main theorem follows immediately from this proposition:

Proof of Theorem 1.3, conditions are sufficient. Let f be the family of pointed stable differentials given by Proposition 4.8. Note that this proposition constructs everything required in Theorem 1.3. In particular, η determines the limit location of the marked zeros and poles, and the components of η are just multiples of the differentials in the degenerating family. It follows that the central fiber of f is indeed $(X, \omega, z_1, \dots, z_n)$. \square

In order to prove Proposition 4.8, we apply induction on the number of levels of the graph. We first use classical plumbing as a base case, then apply higher order plumbing to smooth the nodes connecting the bottom to upper levels, and finally merge dispersed zeros to make sure that we end up in the desired stratum.

Proof of Proposition 4.8. For the induction it will be useful that we allow the curve X to be possibly disconnected. Hence we extend the notion of the number of levels of a graph to the disconnected case as the maximum of numbers of levels over all connected components. The proof is then by induction on the number N of levels of a possibly disconnected curve X .

The *base case of induction* is when the number of levels of X is equal to one, namely, we have $v_1 \asymp v_2$ for any two irreducible components of X . In this case by conditions (1) and (2) all poles at the nodes are simple poles, of opposite residues on the two sides. Moreover, every component is maximal for the order, and thus the pointed stable differential is simply equal to η . This case follows directly by applying classical plumbing given in Proposition 4.4 to every node one by one, which shows that any such differential is plumbable. This procedure obviously preserves the type of the differentials outside the nodes, and we thus obtain a family in the desired stratum $\mathbb{P}\Omega\mathcal{M}_g(\mu)$.

For the *inductive step*, suppose that for any twisted differential η' on a (possibly disconnected) nodal curve X' with a full order $\overline{\Gamma}$ such that the number of levels in $\overline{\Gamma}$ is at most $N - 1$, satisfying all the conditions of the theorem, there exists a degenerating family $(f' : \mathcal{X}' \rightarrow \Delta, \mathcal{W}', \mathcal{Z}_1, \dots, \mathcal{Z}_{n'})$ of pointed stable differentials such that η' is the collection of scaling limits of this family in the sense of Lemma 4.1. Moreover, we may suppose that $\overline{\Gamma}$ is the level graph defined by the function $\ell(\cdot)$ given by the scaling parameters of the family.

To prove Proposition 4.8 by induction, starting with a twisted differential η on X compatible with a full order $\overline{\Gamma}$ with N levels given by a level function ℓ , we need to construct a family degenerating to it, satisfying the conditions claimed. Let L be the minimum value of ℓ on Γ corresponding to the bottom level of $\overline{\Gamma}$. Let $Y_{>L}$ and Z be the (possibly disconnected) stable subcurves of X corresponding to the graphs $\overline{\Gamma}_{>L}$ and $\overline{\Gamma}_{=L}$ respectively.

The restriction of η to $Y_{>L}$ is again a twisted differential, in the sense that it is a twisted differential in our original definition for each connected component of $Y_{>L}$. Since the conditions (0)–(3) are local and condition (4) is imposed level-by-level, we conclude that the restriction of η to $Y_{>L}$ satisfies all the conditions (0)–(4). By the

inductive assumption, there thus exists a family of differentials

$$(f_{>L} : \mathcal{Y} \rightarrow \Delta, \mathcal{W}_{>L} : \Delta^* \rightarrow (f_{>L})_* \omega_{\mathcal{Y}/\Delta}(-\mathcal{Z}_{\text{pol}}))$$

satisfying the conditions in Proposition 4.8.

Next, we want to plumb the family $(\mathcal{Y}, \mathcal{W}_{>L})$ to the constant family $\mathcal{Z} := Z \times \Delta \rightarrow \Delta$ with differential $\eta|_Z$ along the nodes where $\bar{\Gamma}_{>L}$ and $\bar{\Gamma}_{=L}$ intersect, by using higher order plumbing provided by Theorem 4.5. In this theorem the exponent b in (4.7) needs to be divisible by $k+1$. Hence our first task is to adjust the initially chosen level function ℓ in order to satisfy the divisibility constraint.

Let $\{q_j\}_{j \in J}$ be the set of nodes where $\bar{\Gamma}_{>L}$ and $\bar{\Gamma}_{=L}$ intersect. Denote by $k_j \geq 0$ the order of $\mathcal{W}_{>L}$ at the branch q_j^+ of such a node. For convenience we introduce $\kappa_j := k_j + 1$ and let $K := \text{lcm}(\kappa_j : j \in J)$. Pulling back the family $f_{>L} : \mathcal{Y} \rightarrow \Delta$ via the base change $t \rightarrow t^K$ on Δ we can from now on suppose that all the scaling parameters appearing in the family $f_{>L}$ are multiples of K , while the maximum of the scaling parameters is still zero. For ease of notation, we continue to use $\ell(\cdot)$ for the new level function on $\bar{\Gamma}_{>L}$ given by the scaling parameters of $f_{>L}$ after the base change. Because of the choice of K we can find a negative integer as the new value of the lowest level $\ell(Z)$, still denoted by L (again for ease of notation), such that

$$(4.8) \quad \ell(v^+(q_j)) - a_j \kappa_j = L \quad \text{for all } j \in J$$

for some positive integers a_j . In this way, the corresponding level graph remains the same as $\bar{\Gamma}$.

The second task is to construct the modification differential ξ needed in Theorem 4.5. We consider each connected component \mathcal{Y}^0 of \mathcal{Y} separately. We want to define a family of meromorphic differentials $\xi|_{\mathcal{Y}^0}$ such that

$$\text{Res}_{q_j^+}(\xi|_{Y_t^0}) = -\text{Res}_{q_j^-}(\eta|_Z) \quad \text{for all nodes } q_j \text{ in each fiber } Y_t^0 \text{ of } \mathcal{Y}^0.$$

Such a family of differentials exists by Lemma 4.6 under the assumption that the sum of residues at those nodes is zero in each Y_t^0 . If Y_t^0 does not contain a marked point corresponding to a prescribed pole, the global residue condition (4) that we imposed gives precisely the desired assumption. If Y_t^0 contains a marked pole z_p , then we allow $\xi|_{Y_t^0}$ to have a simple pole at z_p as well, with $\text{Res}_{z_p}(\xi|_{Y_t^0})$ equal to minus the sum of all $\text{Res}_{q_j^+}(\xi|_{Y_t^0})$ appearing in the above. In this way, we can still apply Lemma 4.6 to obtain the desired ξ . The modification differential ξ will disappear in the scaling limit, because it is further scaled by a positive power t^b in the setting of Theorem 4.5. Moreover for t small enough, adding $t^b \xi$ does not change the type of the polar part of the differential on the smooth locus of Y_t^0 .

We now plumb the family $\mathcal{Y} \cup \mathcal{Z}$ joined along the nodes q_j , for $j \in J$, where \mathcal{Y} and \mathcal{Z} play the role of \mathcal{X}^+ and \mathcal{X}^- in the notation of Theorem 4.5. For each q_j , set $c = -\ell(v^+(q_j))$ and $b = a_j \kappa_j$ in the notation of Theorem 4.5. Then $b + c = -L$ by (4.8), hence the scaling factor $t^{b+c} = t^{-L}$ for the modification differential ξ and for the differential $\eta|_Z$ on \mathcal{Z} is independent of q_j for $j \in J$. Therefore, we can apply Theorem 4.5 simultaneously to plumb all the nodes q_j , for $j \in J$, with fixtures \mathbb{V}_{a_j} and desired model differentials Ω_j locally around all q_j . As a result, we thus obtain the

plumbed family

$$(f_{\text{nod}} : \mathcal{X}_{\text{nod}} \rightarrow \Delta, \mathcal{W}_{\text{nod}} : \Delta^* \rightarrow (f_{\text{nod}})_* \omega_{\mathcal{X}_{\text{nod}}/\Delta}(-\mathcal{Z}_{\text{pol}}))$$

of differentials whose scaling limit is the twisted differential η and where the condition of non-vanishing on top level still holds.

There are two things left to modify in order to obtain the desired family. First, we apply classical plumbing in Proposition 4.4 to smooth the nodes with simple poles in f_{nod} , originally from the constant family \mathcal{Z} , to obtain a family of differentials

$$(f_{\text{zeros}} : \mathcal{X}_{\text{zeros}} \rightarrow \Delta, \mathcal{W}_{\text{zeros}} : \Delta^* \rightarrow (f_{\text{zeros}})_* \omega_{\mathcal{X}_{\text{zeros}}/\Delta}(-\mathcal{Z}_{\text{pol}}))$$

with smooth fibers for $t \neq 0$, and with the same scaling limit η as that of \mathcal{W}_{nod} . Next, this family $(f_{\text{zeros}}, \mathcal{W}_{\text{zeros}})$ may not belong to the desired stratum, because the orders of marked zeros may have been altered when adding the modification differential ξ to the family of differentials $\mathcal{W}_{>L}$ on \mathcal{Y} . But by Lemma 4.7 there exists a merging in the neighborhood of each marked zero, such that the nearby dispersed zeroes of $\mathcal{W}_{\text{zeros}}$ with total multiplicity m_i are merged back together to form a single zero of multiplicity m_i . We thus finally obtain a family of smooth Riemann surfaces along with differentials in the desired stratum, converging to the given stable curve, and the scaling limit of the differentials is the twisted differential η that we started with. \square

5. FLAT GEOMETRIC SMOOTHING

In this section we give an alternative proof of the sufficiency of the conditions in Theorem 1.3 using techniques from flat geometry. The core ideas of the proof and the induction procedure are parallel to the proof by plumbing in the preceding section. However, the flat geometric pictures presented in this section might look more familiar to some readers. In particular, they make the necessity of the global residue condition quite transparent. One difference is that we construct a degenerating family only over a real segment $[0, \varepsilon)$ rather than a disc Δ_ε as in the proof via plumbing.

For any twisted differential η compatible with a full order, our goal is to construct a family of flat surfaces in the appropriate stratum that makes the degeneration to η visible. We will first describe the construction, and then in the last part of this section justify that the family we construct indeed converges to the desired limit. Below we start with two examples that illustrate the main features of the flat geometric proof.

5.1. Two illustrative examples. In the first example, suppose X is a nodal curve with two irreducible components Y and Z of genus 1 and 2 respectively, joined at two nodes. Let $\bar{\Gamma}$ be a level graph of X such that Y and Z are of the same level. Let $\eta = (\eta_Y, \eta_Z)$ be a twisted differential in terms of the flat geometric pictures on the left and in the middle of Figure 13. In particular, η_Y has two simple poles at q_1^+ and q_2^+ , η_Z has two simple poles at q_1^- and q_2^- , and $\text{Res}_{q_i^+}(\eta_Y) + \text{Res}_{q_i^-}(\eta_Z) = 0$ for $i = 1, 2$, since the widths of the strips agree. The pairs of points (q_i^+, q_i^-) , $i = 1, 2$ at the ends of the infinitely long strips are then glued together to form the two nodes of X . Obviously η is compatible with $\bar{\Gamma}$.

We want to show that the pointed stable differential defined by $\omega = \eta$ with marked points z_i being the zeros of ω is a limit as flat surfaces in the stratum $\Omega\mathcal{M}_4(3, 2, 1)$




FIGURE 13. Smoothing two pairs of simple poles

degenerate. For this purpose, it suffices to truncate each of the half-infinite cylinders at a finite height $R = 1/|t|$ and glue them along 'seam' strips of some fixed height ε , where the *matching residues* at each pair of simple poles ensure that the gluing procedure can be carried out by translation. For $t \rightarrow 0$ the flat surfaces obtained in this construction visibly degenerate to $(X, \omega, z_1, z_2, z_3)$, and we will justify the convergence in general in the proof at the end of the section. This example is an instance of components joined by *flat cylinders* in the language of [Raf07] or, equivalently, the flat geometric viewpoint of classical plumbing (Proposition 4.4).

In the second example, suppose X is a nodal curve with two irreducible components Y and Z of genus 2 and 1 respectively, joined at three nodes. Let $\bar{\Gamma}$ be a level graph of X such that $Y \succ Z$. Let $\eta = (\eta_Y, \eta_Z)$ be a twisted differential in terms of the flat geometric pictures in Figure 14 (without the slit in the interior) and Figure 15. The flat surface Z consists of 5 copies of the Euclidean plane (with modifications in the center), each of which is drawn in Figure 15 as a disk of large radius. Parallel edges with the same labels are identified via translation. We define q_1^- to be the point at infinity of the first two top discs, q_3^- at infinity of the third, and q_2^- at infinite of the bottom two discs. The pairs of points (q_i^+, q_i^-) for $i = 1, 2, 3$ are identified together to form the three nodes of X .

Since the orders of η at q_i^+ and q_i^- add up to -2 and the global residue condition automatically holds by the residue theorem on Z , we conclude that η is a twisted differential compatible with $\bar{\Gamma}$. The stable differential ω associated to η is equal to η_Y on Y , and is identically zero on Z . Denoting by z the unique zero of η_Z , we want to show that (X, ω, z) is a limit of flat surfaces in $\Omega\mathcal{M}_5(8)$, by constructing a family of flat surfaces in this stratum which visibly degenerates to (X, ω, z) .




FIGURE 14. Top level flat surface Y inside $\Omega\mathcal{M}_2(1, 1, 0)$




FIGURE 15. Bottom level flat surface Z inside $\Omega\mathcal{M}_1(8, -2, -3, -3)$

Our strategy is to remove from Y a small disk (as a union of metric half-disks) around each point q_i^+ , scale η_Z by a smaller factor, take a disk of the same size under the flat geometric presentation for the pole q_i^- in the rescaled surface Z , and glue it into Y along the annuli of the same size around q_i^\pm . More precisely, we want to glue along the annuli A_i^+ to A_i^- for $i = 1, 2, 3$, as presented in Figures 14 and 15. To first approximation the total angles around q_i^+ and q_i^- match, because the orders of η_Y and

η_Z add up to -2 at the two branches of a node. The issue is that we cannot directly glue their boundaries, because in each annulus around q_i^- a slit of 'size' of the residue at that pole is missing. In order to remedy this problem, we slit the flat surface Y appropriately, as drawn in Figure 14. The global residue condition ensures that we can remove a central polygon (given by the dashed triangle in this example) from Y , and slit from the points q_i^+ to get a surface in which we can glue to Z along the modified annuli A_i^+ with A_i^- .

We next make the degeneration process more precise by specifying the sizes of the annuli and the slits. Fix $0 < \delta < 1$ close to one. For $t \in (0, \varepsilon)$ sufficiently small, we may assume that the circles of radius $t^{1/2}$ under the flat metric of (Y, η_Y) around the points q_i^+ are disjoint and contain no other special points inside (except for q_i^+). Moreover, we may assume that the circles of radius $t^{1/2}\delta$ under the flat metric of $(Z, t\eta_Z)$ (i.e., rescale Z by t) contain all the interior edges of (Z, η_Z) (those with labels w_i), because these circles have large radius $t^{-1/2}\delta$ under the original flat metric of η_Z .

After these preparations, we can glue the annuli (modified by the neighborhoods of size tr_i around the slits in Y) between the circles of radius $t^{1/2}$ and $t^{1/2}\delta$ with labels A_i^+ to those with labels A_i^- . In particular as $t \rightarrow 0$, the annuli are shrinking to q_i^+ on (Y, η_Y) and expanding to q_i^- on (Z, η_Z) . This gives the desired family converging to (X, ω, z) . Again, we will justify the convergence later in the proof. This example is an instance of components joined by *expanding annuli* in the language of [Raf07], or, equivalently the flat geometric viewpoint of higher order plumbing (Theorem 4.5).

5.2. Construction of degenerating families in the general case. The first step is to prove that a slit as in the second example above exists in general and carry out the construction inductively with respect to a given level graph.

For simplicity of exposition, we restrict to the case of strata of holomorphic type. The same method works for the case of strata of meromorphic type. There is only one place where the global residue condition plays a role to distinguish the two cases, and we will remark on it in the construction below. In addition, we may assume that the level graph does not have an edge joining two vertices on the same level, or equivalently, that a compatible twisted differential has no simple pole at any node. This is because smoothing such a simple polar node with matching residues is a local procedure ensured by classical plumbing, which can be performed in the flat geometric picture according to the first example above.

Let $(X, \omega, q_1, \dots, q_N)$ be a Riemann surface with a meromorphic differential ω and marked points q_1, \dots, q_N that form a subset of zeros and ordinary points of ω . We let X^0 be the open Riemann surface with disks of radius δR removed around the points q_i , where $0 < \delta < 1$ is chosen sufficiently close to one. We refer to the annuli between the circles of radius δR and R as *boundary annuli*.

We next work towards the definition of a residue slit, i.e, a collection of broken lines in the surface around which we will modify neighborhoods of size given by residues. We remark that a residue slit is in general more complicated than just a slit. Given a tuple of sufficiently small complex numbers $T = (r_1, \dots, r_N)$ with $\sum_{i=1}^N r_i = 0$, we fix a permutation $\pi \in S_N$ such that the slopes of $r_{\pi(1)}, \dots, r_{\pi(N)}$ are monotone on S^1 . Let $P = P(T, \pi)$ be the polygon whose edges are given by the vectors $r_{\pi(1)}, \dots, r_{\pi(N)}$

consecutively. It follows that $P(T, \pi)$ is convex. Let $B(P)$ be the barycenter of P . Let p be a point in X^0 , disjoint from the zeros and poles of ω . We place P inside X^0 with $B(P) = p$.

A *residue slit* for (T, π) is then defined to be a collection of broken lines (b_1, \dots, b_n) with the following properties:

- Each broken line b_i starts with the segment from p to the midpoint of the edge $e_{\pi^{-1}(i)}$ of the polygon $P(T, \pi)$, and then connects to q_i . We denote by b_{ij} the line segments constituting b_i and by $\theta(b_{ij})$ the slopes of b_{ij} .
- The broken lines b_i do not intersect (except for the starting point p) and they are disjoint from the zeros and poles of ω (except for the endpoints q_i).
- The slopes $\theta(b_{ij})$ are different from that of $\pm r_i$ for each i .

For $t \in (0, \varepsilon)$ sufficiently small, let $tT = (tr_1, \dots, tr_N)$. We define the surface $X_t^0 = X_t^0(T, \pi)$ obtained by modifying neighborhoods of size tT around the residue slit as follows. Remove $P(T, \pi)$ from X^0 . In the case that $\langle \theta(b_{ij}), r_i \rangle > 0$ (as in Figure 14 for all three broken lines), we *remove* neighborhood parallelograms swept out by planar segments with holonomy vector tr_i centered at points of b_{ij} in general, and glue the parallel sides of these removed parallelograms in pairs. On the other hand for each of the segments where $\langle \theta(b_{ij}), r_i \rangle < 0$, we *add* a parallelogram swept out by segments of holonomy tr_i to the existing flat surface and glue the pieces as indicated in Figure 16. A small modification one needs to carry out is that near a turning point, e.g., the right endpoint of a_3 or the left endpoint of d_1 in Figure 16, we move the removed parallelogram region up or down by $tr_i/2$, which ensures that adding a parallelogram works near the turning point as the residue slit changes its direction. The size of t is constrained by the requirement that the neighborhoods of the residue slit used in this construction are disjoint and do not contain the zeros and poles of ω on X^0 . In this way we obtain the desired surface X_t^0 .




FIGURE 16. A piece of a residue slit and the gluing procedure in the case of a backwards slope

Lemma 5.1. *Given a pointed flat surface $(X, \omega, q_1, \dots, q_N; p)$, for any tuple T of sufficiently small complex numbers with sum equal to zero, there exists a residue slit.*

Proof. We produce the broken lines inductively. At each step we need to find a path from q_i to p , avoiding finitely many contractible subsets, namely the zeros and poles of ω on X^0 and the union of broken lines constructed in the preceding steps. Such a path clearly exists and can be straightened into broken lines, avoiding slopes of $\pm r_i$. \square

Next, we show that every meromorphic differential (without simple poles) can be presented similarly to Figure 15. The building blocks of this construction are basic domains, as introduced by Boissy (see [Boi15a, Section 3.3]). Compared to his definition we ignore infinite cylinders (i.e., simple poles) because such nodes can be smoothed out locally by the classical plumbing, but we need to allow more flexibility in the direction that we slit. Consequently we define for a fixed angle θ a *basic domain in the direction*




FIGURE 17. Two basic domains in the direction $\theta = \pi/4$

θ to be a half-plane with broken polygonal boundary, such that the infinite boundary rays have direction $\pm e^{i\theta}$ and the finite edges w_j of the boundary satisfy $\langle w_j, e^{i\theta} \rangle > 0$ as pictured in Figure 17.

Lemma 5.2. *Let (X, ω) be a meromorphic differential without simple poles, but with l poles of higher order $|m_{n-l+1}|, \dots, |m_n|$, respectively. Then for almost every direction θ there exist $2 \sum_{i=1}^l (|m_{n-l+i}| - 1)$ basic domains in the direction θ such that (X, ω) is obtained by gluing the boundary segments of the basic domains by translation. Moreover, the infinite boundary rays are glued in a way such that the set of basic domains is partitioned in l cycles of lengths $2|m_{n-l+1}| - 2, \dots, 2|m_n| - 2$, respectively.*

If (X, ω) can be represented by basic domains in the direction θ , then all flat surfaces in a neighborhood of (X, ω) also have this property.

Proof. This lemma is a restatement of the infinite zippered rectangle construction of Boissy ([Boi15a]). \square

We are ready to construct the desired family of flat surfaces by induction on the number of levels. Let $(X, \omega, z_1, \dots, z_n)$ be a pointed stable differential associated with a twisted differential η of type μ , where η is compatible with a full order $\bar{\Gamma}$ on X . In the sequel we fix a number $0 < \delta < 1$ sufficiently close to 1.

First consider the case when $\bar{\Gamma}$ has exactly two levels. We want to glue the two levels as in the second example above. For each connected component Y_j of top level, let q_1, \dots, q_N be the nodes where Y_j is joined to the lower level. For $t \in (0, \varepsilon)$ sufficiently small and a positive integer k , we construct residue slits using Lemma 5.1 for $(t^k T_j, \pi)$, where the residues r_i in T_j are given by $r_i = \text{Res}_{q_i^-}(\eta_{v^-(q_i)})$, i.e., they arise from the

lower level branches at the nodes q_i , and the permutation π rearranges the residues in T_j such that their slopes are in some monotone order. Then we add or remove parallelogram neighborhoods of size $t^k T_j$ around the residue slits on the open flat surfaces $(Y_j^0, \eta_{Y_j^0})$, where Y_j^0 is obtained by removing discs of radius $t^{k-1/2}\delta$ around all q_i^+ from (Y_j, η_{Y_j}) with the induced flat metric. As a result, we obtain a continuous family of open flat surfaces (X_t^0, ω_t^0) over $(0, \varepsilon)$, where the removed discs are arbitrarily small as $t \rightarrow 0$.

Next we scale the subsurfaces (Z_j, η_{Z_j}) of lower level as $(Z_j, t^k \eta_{Z_j})$ and present them using Lemma 5.2 for any θ satisfying the conclusion of this lemma. For t small enough, we can assume that the union of *inscribed half-discs* of radius $t^{k-1/2}$ centered at the common limit point as $t \rightarrow 0$ of all broken line segments contains all finite boundary segments in the basic domain presentation for each pole q_i^- in $(Z_j, t^k \eta_{Z_j})$. We then glue these inscribed half-discs into (X_t^0, ω_t^0) , by identifying the boundary annuli A_i^- of inner radius $t^{k-1/2}\delta$ and outer radius $t^{k-1/2}$ with the boundary annuli A_i^+ of the same size (modified by the residue slit construction) for all q_i . We have thus finished the construction for the case of two levels.

Now for general $\bar{\Gamma}$, the induction step follows from the same idea. Let $\ell(\cdot)$ be any integral valued level function on Γ that gives the full order $\bar{\Gamma}$, mapping the top level to zero. Suppose that we have inductively constructed families of flat surfaces $(Y_j(t), \omega_j(t))$ for each of the connected components of $\bar{\Gamma}_{>L}$ for some level L . For given index j , let q_1, \dots, q_N be the nodes joining Y_j to level L . If Y_j does not contain any marked point as a prescribed pole, then the corresponding tuple T_j of residues at all q_i^- has sum equal to zero by the global residue condition. Here is the only place where the global residue condition makes a difference if Y_j contains a marked pole. In that case, there is no global residue condition imposed on it, but we can leverage on the flat geometric presentation of the marked pole to create a residue r_0 such that $r_0 = -\sum_{i=1}^N r_N$. Then the construction as in Figure 14 still goes through by taking the residue tuple $T_j = (r_0, r_1, \dots, r_N)$.

Next we construct a residue slit in the family $(Y_j(t), \omega_j(t))$ as in Lemma 5.1, which has been stated for a single flat surface. A closer look at the proof of that lemma reveals that the changes of complex structures happen in some neighborhoods of the residue slits constructed in the previous inductive steps. Since these finitely many contractible neighborhoods are disjoint from the points q_1^+, \dots, q_N^+ on $Y_j(t)$, the method of the proof still applies in families. Finally we scale the subsurfaces $(Z_v, \eta|_{Z_v})$ with level $\ell(v) = L$ as $(Z_v, t^{-L} \eta_{Z_v})$, present them as in Lemma 5.2, remove disks of radius $t^{-L-1/2}\delta$ around q_i^+ from $Y_j(t)$, and glue in the inscribed half-discs of radius $t^{-L-1/2}\delta$ in $(Z_v, t^{-L} \eta_{Z_v})$ along the boundary annuli of inner radius $t^{-L-1/2}\delta$ and outer radius $t^{-L-1/2}$ (modified by the residue slits of size $t^{-L} T_j$ in $Y_j(t)$). The term $1/2$ in the exponents of t ensures that the annuli are expanding on the original surface (Z_v, η_{Z_v}) and shrinking on $(Y_j(t), \omega_j(t))$ as $t \rightarrow 0$. This thus completes the construction for the general case. As a result, we have constructed a family of flat surfaces (X_t, ω_t) , which is continuous over $(0, \varepsilon)$ and lies in the given stratum of type μ .

Remark 5.3. The flat geometric construction is over a real ray, i.e., the base parameter t is real. It is natural to ask whether one can similarly construct such a family that varies holomorphically over a punctured disk. We point out in general there is a *monodromy* obstruction. For instance, recall the construction by gluing Figure 15 into Figure 14 to form a nearby flat surface in the resulting family. If t varies as a complex parameter, it means that the residue cuts r_i have to vary their arguments besides shrinking the lengths. If the corresponding cycle of r_i is not homologous to zero, when its argument turns back, the Picard-Lefschetz formula implies that it affects those periods that have nonzero intersection with r_i , e.g., the periods arising from the boundary of the residue slit in Figure 14 cannot remain unchanged during the entire residue turning process. Consequently one has to vary holomorphically those affected periods to cancel out the excess periods caused by monodromy, which is less visible in terms of flat geometric coordinates. This phenomenon is closely related to extendability of period coordinates to the boundary of the strata compactification. In [BCGGM] we will study this question systematically.

Remark 5.4. The flat geometric construction of gluing higher order poles along boundary annuli with residue slits is analogous to higher order plumbing in Theorem 4.5. In particular, the residue slit construction plays a similar role as the modification differential in the plumbing construction. However, even over a real ray the two constructions in general do not give the same families if the residues are nonzero. To see this, note that the residue slit construction involves various choices, and so does the plumbing construction, especially when invoking Lemma 4.7 to merge the zeros that were dispersed. Hence they can agree only if we make very specific choices in the respective constructions.

5.3. Flat geometric proof of Theorem 1.3: conditions are sufficient. So far we have constructed a family of flat surfaces (X_t, ω_t) that vary continuously with $t \in (0, \varepsilon)$ sufficiently small. The remaining step is to show that it converges as $t \rightarrow 0$ to the pointed stable differential we started with. For this purpose we need a certain topology on $\overline{\mathcal{M}}_{g,n}$ in which we can verify convergence of the family.

Let (X, Z) be a pointed stable curve, where $Z = \{z_1, \dots, z_n\}$ is the set of marked points on X , and let X' denote X minus its nodes, so X' is a disjoint union of irreducible components of X , punctured at the nodes. An *exhaustion* of X is a sequence of subsets $\dots K_{m-1} \subset K_m \subset K_{m+1} \dots$ of X' , whose union is X' . A sequence of pointed stable curves (X_m, Z_m) in $\overline{\mathcal{M}}_{g,n}$ converges to (X, Z) in the *quasiconformal topology*, if for some exhaustion $\{K_m\}$ of X there exists a sequence of maps $f_m: K_m \rightarrow X_m$ that are quasiconformal onto the images, such that f_m respects the marked points and such that the *dilatation* of f_m tends to 1. The quasiconformal topology was introduced by Abikoff ([Abi77]) in the setting of the augmented Teichmüller space. As $\overline{\mathcal{M}}_{g,n}$ is a complex projective variety, it also inherits a standard topology. Hubbard and Koch ([HK14]) established that these two topologies are equivalent.

For each node q of the stable curve X , let $D_q^+(t)$ be the closed disc of radius $t^{\ell(v^+(q)) - \ell(v^-(q)) - 1/2}$ around q^+ on the higher level branch $(X_{v^+(q)}, \eta_{v^+(q)})$ of q (modified in the neighborhood of the residue slit). In the gluing construction of X_t , the boundary of $D_q^+(t)$ becomes a (modified) circle of radius $t^{-1/2}$ on the lower level branch

$(X_{v^-(q)}, \eta_{v^-(q)})$, which separates the basic domain presentation of the pole q^- into two regions. Let $D_q^-(t)$ be the closure of the region that contains q^- . Denote by $\gamma_q(t)$ the common boundary curve of $D_q^\pm(t)$ in X_t after the gluing construction. Denote by $D^\pm(t)$ the union of $D_q^\pm(t)$ and by $\gamma(t)$ the union of $\gamma_q(t)$ over all nodes q .

Let $K_t := X \setminus D^\pm(t)$. Then $\{K_t\}$ is an exhaustion of X as $t \rightarrow 0$. Moreover, K_t looks just alike $X_t \setminus \gamma(t)$, and the only difference is that X_t is further modified around the neighborhoods of the residue slits outside $D^+(t)$. This difference can be measured by the following result, which is the main step towards showing that the family of flat surfaces we have constructed actually converges to the prescribed limit object.

Lemma 5.5. *As $t \rightarrow 0$, there exists a quasiconformal map $f_t: K_t \rightarrow X_t \setminus \gamma(t)$, respecting the marked points $z_i(t)$, such that the dilatation of f_t tends to 1.*

For a simple example, consider a two-level surface with a torus on the top level joined to the lower level at two nodes q_1 and q_2 . Suppose the value of the lower level is -1 and the direction of the residues $\pm r$ at the two nodes is vertical. Then the map f_t is given in Figure 18. Only the top level surface is depicted, since there is no residue slit on the lower level. The residue slit in this case is just a straight line joining q_1^+ to q_2^+ , presented as the dotted line in the picture.




FIGURE 18. The quasiconformal map f_t on the top level

The quasiconformal map f_t in this case is the identity outside the union of the two disks $D_{q_1}^+(t)$, $D_{q_2}^+(t)$, and the rectangular region swept out by segments of holonomy $t^{1/2}(2\rho)$, for some fixed $\rho < 1$, centered around the residue slit. We choose the scaling factor $t^{1/2}$ because it is the radius of the disks $D_{q_i}^+(t)$, hence the rectangular region goes into the disks under the assumption on ρ . Inside the rectangular region (and still outside of the disks), f_t takes the upper (resp. lower) half in the left picture to the upper (resp. lower) half in the right picture, by a family of linear maps preserving vertical lines and shortening the height of the vertical segments by that of the removed residue slit neighborhood, with a small perturbation near the boundary of $D_{q_i}^+(t)$. Note that f_t is continuous along the dotted line on the left, because the upper and lower boundary segments of the residue slit neighborhood are identified in the image surface X_t on the right. Moreover, since the vertical segments are shortened by at most $tr/2$, which becomes arbitrarily small compared to $t^{1/2}\rho$ as $t \rightarrow 0$, it implies that the dilatation of f_t tends to 1.

Proof of Lemma 5.5. The idea of the proof is similar to the example presented above. Let b_{ij} be a broken line segment of the residue slit on the subsurface (X_v, η_v) that was constructed when gluing in a subsurface at level $L < \ell(v)$. If $\langle \theta(b_{ij}), r_i \rangle > 0$, consider the neighborhood of b_{ij} swept out by segments of holonomy $t^{\ell(v)-L-1/2}(2\rho)$ centered around the residue slit. On the upper and lower parts of this neighborhood we define the quasiconformal map f_t to be an affine transformation onto the respective $(t^{\ell(v)-L-1/2}(\rho - t^{1/2}r_i/2))$ -segments as illustrated in Figure 18, and then composed with scaling by $t^{-\ell(v)}$, because (X_v, η_v) is scaled by $t^{-\ell(v)}$ before gluing into X_t . If $\langle \theta(b_{ij}), r_i \rangle < 0$ we use the same composition, except that the image of the affine transformation in the upper and lower parts will be swept out by the $(t^{\ell(v)-L-1/2}(\rho + t^{1/2}r_i/2))$ -segments. Since the ratio of ρ and $\rho \pm t^{1/2}r_i/2$ tends to 1 as $t \rightarrow 0$, it follows that the dilatation of f_t on these neighborhoods of the residue slit tends to 1.

Moreover, in the neighborhood of a corner where the residue slit changes its direction from $\langle \theta(b_{ij}), r_i \rangle > 0$ to $\langle \theta(b_{ij}), r_i \rangle < 0$, the quasiconformal map f_t is the identity composed with scaling by $t^{-\ell(v)}$. To see this, e.g., take the angular sector between a_3 and b in Figure 16, slide it up so that the sides a_3 touch, and glue in the additional parallelogram. One can easily find linear maps along segments in the direction r_i that interpolate continuously from the situation generically along the broken line to the situation near such a corner.

In the central polygon P , we extend the residue slit by connecting the middle points of the edges to the barycenter $B(P)$. We define the quasiconformal map f_t in a neighborhood of this polygon by affine transformations that stretch along the lines joining the extended slit segments to the respective vertices of the polygon neighborhood as in the figure to the right, and then composed with scaling by $t^{-\ell(v)}$. Again, one can find a continuous family of linear maps along segments whose direction varies from r_i to r_{i+1} near the central polygon. Such a map is sketched in Figure 19, where the dashed lines represent curves along which f_t stretches.




FIGURE 19. The quasiconformal map f_t nearby the central polygon

On the complement of these neighborhoods of the residue slits in each component (X_v, η_v) , we take f_t to be the scaling map by $t^{-\ell(v)}$, which glues continuously to the quasiconformal map on the neighborhoods constructed above.

It is clear from the construction, as explained above, that the dilatation of f_t depends continuously on t and tends to 1 as $t \rightarrow 0$. \square

Lemma 5.5 proves the desired convergence in $\overline{\mathcal{M}}_{g,n}$. Nevertheless, it does not quite imply convergence of pointed stable differentials in the incidence variety compactification, as explained in Remark 2.2. In order to show that (X_t, ω_t) converges to η_v projectively on each component X_v of X , it suffices to show that the location and multiplicity of all zeros and poles agree. This is obvious for all zeros and poles of η in the smooth locus of X . It remains to check the vanishing orders of η at the nodes of X (compared to Example 3.2). At each node q , the vanishing order can be detected by the index of a path centered around q^+ under the flat metric in the higher level branch of q . Since this quantity is defined in a neighborhood of q^+ , untouched in the construction for t small enough, the vanishing orders of the limit of ω_t as $t \rightarrow 0$ are equal to those of η at every node.

To show that (X_t, ω_t) converges not only projectively, but also with the given scales at top level, it suffices to compare ratios of the length of a nonzero (relative or absolute) period in each top level component. This is obvious for a top level component of positive genus or with more than one marked zero. In the remaining cases, such a component has at least two nodes q_1 and q_2 joining it to lower level. A path on X_t joining a boundary point on each annulus A_i^+ around q_i (that we used for gluing X_t) converges to a path joining q_1 to q_2 , and hence can be used for comparison of relative size. \square

REFERENCES

- [Abi77] W. Abikoff. *Degenerating families of Riemann surfaces*. Ann. Math. (2) 105.1 (1977), pp. 29–44.
- [AM69] M. Atiyah and I. Macdonald. *Introduction to commutative algebra*. Reading, Mass.-London-Don Mills, Ont: Addison-Wesley Publishing Co., 1969, pp. ix+128.
- [Bai07] M. Bainbridge. *Euler characteristics of Teichmüller curves in genus two*. Geom. Topol. 11 (2007), pp. 1887–2073.
- [BCGGM] M. Bainbridge, D. Chen, Q. Gendron, S. Grushevsky, and M. Möller. *Boundary structure of strata of abelian differentials*. In preparation.
- [Boi15a] C. Boissy. *Connected components of the strata of the moduli space of meromorphic differentials*. Comment. Math. Helv. 90.2 (2015), pp. 255–286.
- [Boi15b] C. Boissy. *Moduli space of meromorphic differentials with marked horizontal separatrices* (2015). arXiv: 1507.00555.
- [CC18] D. Chen and Q. Chen. *Principal boundary of moduli spaces of abelian and quadratic differentials*. Ann. Inst. Fourier (2018, to appear).
- [Che17] D. Chen. *Degenerations of Abelian differentials*. J. Differential Geom. 107.3 (2017), pp. 395–453.
- [EH86] D. Eisenbud and J. Harris. *Limit linear series: basic theory*. Invent. Math. 85.2 (1986), pp. 337–371.
- [EH87] D. Eisenbud and J. Harris. *Existence, decomposition, and limits of certain Weierstrass points*. Invent. Math. 87.3 (1987), pp. 495–515.
- [EKZ14] A. Eskin, M. Kontsevich, and A. Zorich. *Sum of Lyapunov exponents of the Hodge bundle with respect to the Teichmüller geodesic flow*. Publ. Math. Inst. Hautes Études Sci. 120 (2014), pp. 207–333.

- [EM02] E. Esteves and N. Medeiros. *Limit canonical systems on curves with two components*. Invent. Math. 149.2 (2002), pp. 267–338.
- [EM13] A. Eskin and M. Mirzakhani. *Invariant and stationary measures for the $SL(2, \mathbb{R})$ action on Moduli space* (2013). arXiv: 1302.3320.
- [EMM15] A. Eskin, M. Mirzakhani, and A. Mohammadi. *Isolation, equidistribution, and orbit closures for the $SL(2, \mathbb{R})$ action on moduli space*. Ann. Math. (2) 182.2 (2015), pp. 673–721.
- [EMR12] A. Eskin, M. Mirzakhani, and K. Rafi. *Counting closed geodesics in strata* (2012). arXiv: 1206.5574.
- [EMZ03] A. Eskin, H. Masur, and A. Zorich. *Moduli spaces of Abelian differentials: the principal boundary, counting problems, and the Siegel-Veech constants*. Publ. Math. Inst. Hautes Études Sci. 97 (2003), pp. 61–179.
- [Fil16] S. Filip. *Splitting mixed Hodge structures over affine invariant manifolds*. Ann. Math. (2) 183.2 (2016), pp. 681–713.
- [For91] O. Forster. *Lectures on Riemann surfaces*. Vol. 81. Graduate Texts in Mathematics. Springer-Verlag, New York, 1991, pp. viii+254.
- [FP16] G. Farkas and R. Pandharipande. *The moduli space of twisted canonical divisors*. J. Inst. Math. Jussieu (2016, to appear).
- [Gen18] Q. Gendron. *The Deligne-Mumford and the Incidence Variety Compactifications of the Strata of $\Omega\mathcal{M}_g$* . Ann. Inst. Fourier (2018, to appear).
- [GKN17] S. Grushevsky, I. Krichever, and C. Norton. *Real-normalized differentials: limits on stable curves* (2017). arXiv: 1703.07806.
- [HK14] J. Hubbard and S. Koch. *An analytic construction of the Deligne-Mumford compactification of the moduli space of curves*. J. Differential Geom. 98.2 (2014), pp. 261–313.
- [Hu17] X. Hu. *The locus of plane quartics with a hyperflex*. Proc. Amer. Math. Soc. 145.4 (2017), pp. 1399–1413.
- [KZ03] M. Kontsevich and A. Zorich. *Connected components of the moduli spaces of Abelian differentials with prescribed singularities*. Invent. Math. 153.3 (2003), pp. 631–678.
- [Lan93] S. Lang. *Real and functional analysis*. Third. Vol. 142. Graduate Texts in Mathematics. Springer-Verlag, New York, 1993, pp. xiv+580.
- [LU17] B. Lin and M. Ulirsch. “Towards a tropical Hodge bundle”. In: *Smith G., Sturmfels B. (eds) Combinatorial Algebraic Geometry*. Vol. 80. Fields Institute Communications. Springer, New York, NY, 2017, pp. 353–368.
- [Mul17] S. Mullane. *On the effective cone of $\overline{\mathcal{M}}_{g,n}$* . Adv. Math. 320 (2017), pp. 500–519.
- [MUW17] M. Möller, M. Ulirsch, and A. Werner. *Realizability of tropical canonical divisors* (2017). arXiv: 1710.06401.
- [MW17] M. Mirzakhani and A. Wright. *The boundary of an affine invariant submanifold*. Invent. Math. 209.3 (2017), pp. 927–984.
- [Raf07] K. Rafi. *Thick-thin decomposition for quadratic differentials*. Math. Res. Lett. 14.2 (2007), pp. 333–341.
- [Sau17] A. Sauvaget. *Cohomology classes of strata of differentials* (2017). arXiv: 1701.07867.

- [Str84] K. Strebel. *Quadratic differentials*. Vol. 5. Ergebnisse der Mathematik und ihrer Grenzgebiete (3). Springer-Verlag, Berlin, 1984, pp. xii+184.
- [Whi65] E. F. Whittlesey. *Analytic functions in Banach spaces*. Proc. Amer. Math. Soc. 16 (1965), pp. 1077–1083.
- [Wol13] S. Wolpert. *Infinitesimal deformations of nodal stable curves*. Adv. Math. 244 (2013), pp. 413–440.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405, USA
E-mail address: `mabainbr@indiana.edu`

DEPARTMENT OF MATHEMATICS, BOSTON COLLEGE, CHESTNUT HILL, MA 02467, USA
E-mail address: `dawei.chen@bc.edu`

INSTITUT FÜR ALGEBRAISCHE GEOMETRIE, LEIBNIZ UNIVERSITÄT HANNOVER, WELFENGARTEN 1,
30167 HANNOVER, GERMANY
Current address: Centro de Ciencias Matemáticas-UNAM, Antigua Car. a Pátzcuaro 8701, Col.
Ex Hacienda San José de la Huerta, Morelia, Mich., México
E-mail address: `gendron@matmor.unam.mx`

MATHEMATICS DEPARTMENT, STONY BROOK UNIVERSITY, STONY BROOK, NY 11794-3651, USA
E-mail address: `sam@math.stonybrook.edu`

INSTITUT FÜR MATHEMATIK, GOETHE-UNIVERSITÄT FRANKFURT, ROBERT-MAYER-STR. 6-8, 60325
FRANKFURT AM MAIN, GERMANY
E-mail address: `moeller@math.uni-frankfurt.de`